IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

the application of: KUSTERS ET AL.

Serial Number: 09/904,994

Group Art Unit: To be assigned

Filed: July 13, 2001

Examiner: To be assigned

For: HELICOBACTER FELIS VACCINE

CLAIM TO PRIORITY UNDER 35 USC 119

Assistant Commissioner of Patents Washington, D.C. 20231

Attorney Docket No. 2000.566 US

Akzo Nobel Patent Department

(301) 948-7400

(301) 948-9751

1300 Piccard Drive, Suite 206

Rockville, Maryland 20850-4373

July 30, 2001

03CU #4

Sir:

The benefit of the filing date of the following prior foreign application is hereby requested for the above-identified application, and the priority provided in 35 USC 119 is hereby claimed:

European Patent Application No. 00202565.8, filed July 17, 2000

In support of this claim, the requisite certified copy of said original foreign application is filed herewith along with a verified English translation thereof.

It is requested that the file of this application be marked to indicate that the Applicant has complied with the requirements of 35 USC 119 and that the Patent and Trademark Office kindly acknowledge receipt of this document.

In the event any fees are required with this paper, please charge our Deposit Account No. 02-2334.

Respectfully submitted,

William M. Blackstone Attorney for Applicants Registration No. 29,772

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Assistant Commissioner of Patents, Washington, D.C. 20231 on 7/30/0 20231 on .

Date Of Deposit

71 KUSTERS CLAIMTOPRIORITY

Registration No. 29,772

Signatura

WMB:lcf

Tel:

Fax:

Enclosures

•• :



Europäisches Patentamt

Eur pean Patent Office

Office eur péen des brevets

Blatt 2 der Bescheinigung Sheet 2 of the certificate Page 2 de l'attestation

BEST AVAILABLE COPY

Anmeldung Nr.: Application no.: Demande n*:

00202565.8

Anmeldetag: Date of filing: Date de dépôt:

17/07/00

Anmelder: Applicant(s): Demandeur(s):

Demandeur(s): Akzo Nobel N.V. 6824 BM Arnhem NETHERLANDS

Bezeichnung der Erfindung: Title of the invention: Titre de l'invention:

Helicobacter felis vaccine

In Anspruch genommene Prioriät(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat: State: Pays: Tag: Date: Aktenzeichen: File no. Numéro de dépôt:

Internationale Patentklassifikation: International Patent classification: Classification internationale des brevets:

C12N9/80, C07K14/205, C07K16/12, C12Q1/68, A61K39/106, A61K48/00, G01N33/53, G01N33/68

Am Anmeldetag benannte Vertragstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/TR
Etats contractants désignés lors du depôt:

Bernerkungen: Remarks: Remarques:



European **Patent Office**

Office européen des brevets

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application conformes à la version described on the following page, as originally filed.

Les documents fixés à cette attestation sont initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patent application No. Demande de brevet nº Patentanmeldung Nr.

00202565.8

Der Präsident des Europäischen Patentamts;

For the President of the European Patent Office

Le Président de l'Office européen des brevets

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN THE HAGUE, LA HAYE, LE

10/07/01

,					
	:			-	
			•		·
					•
		·			





EPO - DG DESO

17 07. 2000



Helicobacter felis vaccine.

The present invention relates to novel *Helicobacter* urease subunit polypeptides, nucleic acid sequences encoding these polypeptides, to the polypeptides for use in vaccines and for use in the manufacturing thereof, to vaccines comprising said polypeptides and to methods for the preparation of such vaccines. Further, the invention relates to diagnostic methods for the detection of the nucleic acid sequences, the polypeptides and antibodies against the polypeptides.

- 10 Several Helicobacter species are the cause of pathogenesis of the gastric epithelium. Helicobacter pylori, and to a lesser extent H. heilmannii are known to cause gastritis, a major factor in the development of peptic ulcers and gastric lymphoma in humans. Helicobacter felis is most likely the cause of gastric infections in both cats and dogs. In order to survive the highly acidic environment of the stomach, members of the
- Helicobacter family produce an urease that is capable of hydrolysing the urea present in gastric juice. This hydrolysation sets free an amount of NH₄OH that suffices to neutralise the environment of the bacterium. It is known, that the urease plays a role in the colonisation of the bacterium as well as in its pathogenesis.
- Genes encoding urease have been described and sequenced for both *Helicobacter* pylori (Labigne et al., J. Bacteriol. 173: 1920-1931 (1991)) and *Helicobacter felis* (Ferrero et al., Molec. Microbiol. 9, 323-333 (1993)). Of the seven genes involved in urease expression and secretion, only two genes encode the two structural subunits urease A en B of the urease enzyme; ureA and ureB. These two polypeptides form a polypeptide complex having urease activity.
- Vaccines against infections caused by both *H. pylori* and *felis* have been made and have been the subject of i.a. International Patent Applications WO 94/09823 and WO 96/34624. Several attempt have been made, to use *H. pylori* urease as a vaccine component for the protection of cats against *H. felis* infection. Although indeed a certain level of protection can be obtained, the results are far from the 100 % protection that
- would be desirable. From animal experiments published so far it becomes clear that a significant number of animals vaccinated with *H. pylori* is not at all protected against subsequent challenge with *H. felis*. Protection of cats vaccinated with purified urease from either *H. felis* or *pylori* has not been described. Vaccinating cats with *H. felis* whole cell lysates might theoretically be feasible but is not a practical option. This is because in spite of many attempts for improvement, *H. felis* is difficult to grow.
 - There clearly is a need for an efficacious vaccine, based upon homologous components, and it is clear that the known *H. felis* urease does not confer full protection.
- It is i.a. an object of the present invention to provide a *H. felis* urease which is able to induce protection against *Helicobacter felis* infection in dogs and cats. It was surprisingly found now, that in *H. felis* a second urease exists, of which the genes encoding the structural subunits share only low homology with the known *H. felis* ureA and B genes. The novel urease is named ureaseXY, in order to discriminate it from the known urease AB. The newly found urease has been discovered in *H. felis*, and is not present in *H. pylori*.
 - The overall genetic structure of the genes encoding the two structural urease subunits, UreX and UreY is comparable to that of the known UreA and B in *H. felis* and *H. pylori*. The sequence homology is however surprisingly low. It was "v" n more surprisingly found, that the homology between the ureA and B genes and the novel ureX and Y
- genes in one single *H. felis* strain is even strikingly lower than the homology between the various ureA and B gen s from the various *Helicobacter* species.



Table 1a, 1b and 1c show the comparison of the ureX and Y gene and the polypeptides they encode form five different *Helicobacter felis* species, with the ureA and B genes and polypeptides from *Helicobacter felis*, *pylori* and *heilmannii*.

The level of homology of the genes encoding the novel structural urease subunits X and Y and the polypeptides they encode as compared to that of known ureA and B genes and polypeptide subunits is presented in table 1a, b and c.

10

5

Reference molecule: H. felis ureX CS1	a.a.	n.a.
H. felis ureA	50 %	57 %
H. pylori ureA	52 %	60 %
H. heilmannii ureA	54 %	62 %
H. felis strain Kukka ureX	100 %	91 %
H. felis strain Ds4 ureX	99 %	91 %
H. felis strain 2301 ureX	99 %	91 %
H. felis strain 390 ureX	99 %	91 %

<u>Table 1a:</u> amino acid and nucleic acid homology between the *H. felis* ureX and various ureA subunits.

15

Reference molecule: H. felis ureY CS1	a.a.	n.a.
H. felis ureB	73 %	71 %
H. pylori ureB	73 %	70 %
H. heilmannii ureB	74 %	71 %
H. felis strain Kukka ureY	99 %	95 %
H. felis strain Ds4 ureY	98 %	94 %
H. felis strain 2301 ureY	99 %	95 %

20

<u>Table 1b:</u> amino acid and nucleic acid homology between the *H. felis* ureY and various ureB subunits.

Reference molecule: H. felis ureXY CSI	n.a.
H. felis ureAB	67 %
H. pylori ureAB	67 %
H. heilmannii ureAB	68 %
H. felis strain Kukka ureXY	94 %
H. felis strain Ds4 ureXY	94 %
H. felis strain 2301 ureXY	94 %

25

<u>Table 1c:</u> nucleic acid homology between *H. felis* ureXY and various ureAB genes.







One embodiment of the invention thus relates to nucleic acid sequences encoding the novel urease X and Y subunits.

First of all, this embodiment of the invention relates to nucleic acid sequences encoding two subunits of a urease complex such as expressed by Helicobacter felis, that have at least 85 % homology with SEQ ID NO: 1, or parts thereof with a length of at least 40, preferably 45, more preferably 50 nucleotides encoding at least an immunogenic fragment of one of the subunits. Still even longer fragments, with a length of at least 55, 60 or 70 nucleic acids are in that order even more preferred.

10

30

35

5

A preferred form of this embodiment relates to nucleic acid sequences that encode the urease X subunit polypeptide or the urease Y subunit polypeptide and that have at least 85 % homology with SEQ ID NO: 1, or parts thereof with a length of at least 40. preferably 45, more preferably 50 nucleotides encoding at least an immunogenic fragment of the urease X subunit polypeptide or the urease Y subunit polypeptide.

15 Merely as an example: the nucleic acid sequence encoding the urease X subunit of Helicobacter felis strain CS1 starts at position 206/207/208 (GTG) (See figure 1a (1)) and stops at position 884/885/886 (TAA). the nucleic acid sequence encoding the urease Y subunit of Helicobacter felis strain CS1 starts at position 897/898/899 (ATG) 20 and stops at position 2601/2602/2603 (TAG).

Still even longer fragments, with a length of at least 55, 60 or 70 nucleic acids are in that order even more preferred.

A more preferred form of this embodiment relates to nucleic acid sequences having at 25 least 90 %, preferably 94 %, more preferably 97 % homology therewith.

The determination of the homology percentages was done with the computer program Align Plus for Windows, available from Scientific and Educational Software, P.O.Box 72045 Durham, NC 27722-2045, USA. Settings used for the nucleic acid comparisons are indicated in figures 1a, 1b and 1c.

Since the present invention discloses nucleic acid sequences encoding novel structural Helicobacter felis urease subunits, it is now for the first time possible to obtain such polypeptides in sufficient quantities. This can e.g. be done by using expression systems to express the genes encoding the UreX and UreY subunits. Therefore, in a more preferred embodiment, the invention relates to DNA fragments comprising a nucleic acid sequence according to the invention. Such DNA fragments

can e.g. be plasmids, into which a nucleic acid sequence according to the invention is cloned. Such DNA fragments are useful e.g. for enhancing the amount of DNA for use

40 as a probe, as described below.

> An essential requirement for the expression of the nucleic acid sequence is an adequate promoter operably linked to the nucleic acid sequence. It is obvious to those skilled in the art that the choice of a promoter extends to any eukaryotic, prokaryotic or viral promoter capable of directing gene transcription in cells used as host cells for protein

expression. Therefore, an even more preferred form of this embodiment relates to a recombinant DNA molecule comprising a DNA fragment or a nucleic acid sequence according to the invention that is placed under the control of a functionally linked promotor. This can b

obtained by means of e.g. standard molecular biology techniques. (Maniatis/Sambrook (Sambrook, J. Molecular cloning: a laboratory manual, 1989. ISBN 0-87969-309-6).

Printed:10-07-2001



10

15



4

Functionally linked promotors are promotors that are capable of controlling the transcription of the nucleic acid sequences to which they are linked. When the host cells are bacteria, useful expression control sequences which may be used include the Trp promoter and operator (Goeddel, et al., Nucl. Acids Res., 8, 4057, 1980); the lac promoter and operator (Chang, et al., Nature, 275, 615, 1978); the outer membrane protein promoter (Nakamura, K. and Inouge, M., EMBO J., 1, 771-775, 1982); the bacteriophage lambda promoters and operators (Remaut, E. et al., Nucl. Acids Res., 11, 4677-4688, 1983); the α-amylase (B. subtilis) promoter and operator, termination sequences and other expression enhancement and control sequences compatible with the selected host cell.

When the host cell is yeast, useful expression control sequences include, e.g., α-mating factor. For insect cells the polyhedrin or p10 promoters of baculoviruses can be used (Smith, G.E. et al., Mol. Cell. Biol. 3, 2156-65, 1983). When the host cell is of mammalian origin illustrative useful expression control sequences include the SV-40 promoter (Berman, P.W. et al., Science, 222, 524-527, 1983) or the metallothionein promoter (Brinster, R.L., Nature, 296, 39-42, 1982) or a heat shock promoter (Voellmy et al., Proc. Natl. Acad. Sci. USA, 82, 4949-53, 1985).

Bacterial, yeast, fungal, insect and mammalian cell expression systems are very frequently used systems. Such systems are well-known in the art and generally available, e.g. commercially through Clontech Laboratories, Inc. 4030 Fabian Way, Palo Alto, California 94303-4607, USA. Next to these expression systems, parasite-based expression systems are very attractive expression systems. Such systems are e.g. described in the French Patent Application with Publication number 2 714 074, and in US NTIS Publication No US 08/043109 (Hoffman, S. and Rogers, W.: Public. Date 1 December 1993).

Thus a still even more preferred form of this embodiment of the invention relates to Live Recombinant Carrier micro-organisms (LRCs) comprising a gene encoding the UreX or UreY polypeptide or an immunogenic fragment thereof according to the invention. Such micro-organisms are e.g. bacteria and viruses. These LRC micro-organisms are micro-organisms in which additional genetic information, in this case a gene encoding the UreX or UreY polypeptide or an immunogenic fragment thereof according to the invention has been cloned. Animals infected with such LRCs will produce an immunogenic response not only against the immunogens of the vector, but also against the immunogenic parts of the polypeptide(s) for which the genetic code is additionally cloned into the LRC, e.g. the ureX or Y gene.

As an example of bacterial LRCs, attenuated Salmonella strains known in the art con

As an example of bacterial LRCs, attenuated Salmonella strains known in the art can attractively be used.

Live recombinant carrier parasites have i.a. been described by Vermeulen, A. N. (Int. Journ. Parasitol. 28: 1121-1130 (1998))


Also, LRC viruses may be used as a way of transporting the nucleic acid sequence into a target cell. Live recombinant carrier viruses are also called vector viruses. The site of integration of the gene encoding a UreX or Y polypeptide may be a site in a viral gene that is not essential to the virus, or a site in an intergenic region. Viruses often used as

that is not essential to the virus, or a site in an intergenic region. Viruses often used as vectors are Vaccinia viruses (Panicali et al; Proc. Natl. Acad. Sci. USA, 79: 4927 (1982), Herpesviruses (E.P.A. 0473210A2), and Retroviruses (Valerio, D. et al; in Baum, S.J., Dicke, K.A., Lotzova, E. and Pluznik, D.H. (Eds.), Experimental Haematology today - 1988. Springer Verlag, New York: pp. 92-99 (1989)).

Figure 1a (3

2048) agaatttggtaagcttcctgaagatggcaaagataacttccgcattaagcgctacatctccaaatacactatcaacccgctttgacccacggcgtgagcgagtatatcggctc 1890) tgct.gat.gct.gat.gct.gat.gct.gat.gct.gat.gct.gat.gct.gat.gct.gat.gct.gct.gat.gct.gat.gct.gat.gat.gat.gat.gat.gat.gat.gat.gat.gat.gat.gat.gat.t.t.t	2168) tgtggaagatggccgacttggtggtgtggtatctctttttggcgtaaaacccaaaatcgtgatcaagagggtatggtgtcttctctgaaatggcgattctaacgc 2010) 1964) 1965 1965 1965 1965 1965 1965 1965 1965	2288) gtctgtgcccactccccaaaccggtttattaccgcgaaatgtttgggcatcacggcaaggcgaaatttgacaccagcattactttgtttccaaagtcgcctatgaaaatggcgtgaaaga 2130) 2084) .c. t.a.t.g. 2085) .c. t.a.t.g. 2180) .a.t.g. 2180) .a.t.g.c.g.t.c.a.a.a.a.t.g. 2180) .a.t.c.g.t.c.a.a.a.a.t.g.c.a.a.a.t.a.c.t.t.a.t.a.t.a.c.t.a.c.t.g.a.a.a.tc.g.tc.g.t	2250) a.a.a.ta.g.g.g.c.c.g.g	2528) ctttgtagatggcaaactctgcaactccgcaagtgcctctagcccagcgctacactttcttctag 2370)
Cslseq Kukkaseq (Ds4seq 2301SEQ (H. felis com (H. pylori co (H. heilmannı (Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co (Cslseq Kukkaseq (Kukkaseq (S4seq (Z301SEQ (390seq (H. felis com (H. pylori co (H. pylori co (H. heilmanni)	Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co H. heilmanni	Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co H. heilmanni

Figure 1a (4





The technique of *in vivo* homologous recombination, well-known in the art, can be used to introduce a recombinant nucleic acid sequence into the genome of a bacterium, parasite or virus of choice, capable of inducing expression of the inserted nucleic acid sequence according to the invention in the host animal.

5

10

15

20

25

30

Finally another form of this embodiment of the invention relates to a host cell comprising a nucleic acid sequence encoding a polypeptide according to the invention, a DNA fragment comprising such a nucleic acid sequence or a recombinant DNA molecule comprising such a nucleic acid sequence under the control of a functionally linked promotor. This form also relates to a host cell containing a live recombinant carrier containing a nucleic acid molecule encoding a UreX or Y polypeptide or an immunogenic fragment thereof according to the invention.

A host cell may be a cell of bacterial origin, e.g. Escherichia coli, Bacillus subtilus and Lactobacillus species, in combination with bacteria-based plasmids as pBR322, or bacterial expression vectors as pGEX, or with bacteriophages. The host cell may also be of eukaryotic origin, e.g. yeast-cells in combination with yeast-specific vector molecules, or higher eukaryotic cells like insect cells (Luckow et al; Bio-technology 6: 47-55 (1988)) in combination with vectors or recombinant baculoviruses, plant cells in combination with e.g. Ti-plasmid based vectors or plant viral vectors (Barton, K.A. et al; Cell 32: 1033 (1983), mammalian cells like Hela cells, Chinese Hamster Ovary cells (CHO) or Crandell Feline Kidney-cells, also with appropriate vectors or recombinant viruses.

Another embodiment of the invention relates to the polypeptides encoded by the nucleic acid sequences, i.e. the urease X subunit and the urease Y subunit and to immunogenic fragments thereof according to the invention.

Therefore, this embodiment of the invention relates to the *Helicobacter felis* urease X polypeptide, said polypeptide having an amino acid sequence that is at least 85 % homologous to SEQ ID NO: 2 or an immunogenic fragment of that polypeptide with a length of at least 40 amino acids that is capable of inducing an immune response against ureaseXY. Preferably, the length is more than 40 amino acids, preferably at least 45, 50, 55, 60 or 70 amino acids in that order or preference.

More preferably this embodiment relates to such polypeptides having a sequence homology of at least 90 %, more preferably 95 % homology to SEQ ID NO: 2, or an immunogenic fragment of that polypeptide with a length of at least 40 amino acids that is capable of inducing an immune response against ureaseXY.

This embodiment of the invention also relates to the *Helicobacter felis* urease Y polypeptide, said polypeptide having an amino acid sequence that is at least 85 % homologous to SEQ ID NO: 3 or an immunogenic fragment of that polypeptide with a length of at least 40 amino acids that is capable of inducing an immune response against ureaseXY. Preferably, the length is more than 40 amino acids, preferably at least 45, 50, 55, 60 or 70 amino acids in that order or preference.

45

More preferably this embodiment relates to such polypeptides having a sequence homology of at least 90 %, more preferably 95 % homology to SEQ ID NO: 3, or an immunog nic fragment of that polypeptid with a length of at least 40 amino acids that is capable of inducing an immune response against ureaseXY.



00202335.8(47:07-2000)



6

As for the nucleotide sequence comparison, the comparison between the various amino acid sequences was made using Align Plus for Windows, available from Scientific and Educational Software, P.O.Box 72045 Durham, NC 27722-2045, USA. Settings used for the amino acid comparisons are indicated in figures 1a, 1b and 1c.

5

10

15

20

It will be understood that, for the particular polypeptides embraced herein, natural variations can exist between individual Helicobacter felis strains. These variations may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of (an) amino acid(s) in said sequence. Amino acid substitutions which do not essentially alter biological and immunological activities, have been described, e.g. by Neurath et al in "The Proteins" Academic Press New York (1979). Amino acid replacements between related amino acids or replacements which have occurred frequently in evolution are, inter alia. Ser/Ala, Ser/Gly, Asp/Gly, Asp/Asn, Ile/Val (see Dayhof, M.D., Atlas of protein sequence and structure, Nat. Biomed. Res. Found., Washington D.C., 1978, vol. 5, suppl. 3). Other amino acid substitutions include Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val. Thr/Phe, Ala/Pro, Lys/Arg, Leu/Ile, Leu/Val and Ala/Glu. Based on this information. Lipman and Pearson developed a method for rapid and sensitive protein comparison (Science, 227, 1435-1441, 1985) and determining the functional similarity between homologous proteins. Such amino acid substitutions of the exemplary embodiments of this invention, as well as variations having deletions and/or insertions are within the scope of the invention as long as the resulting polypeptides retain their immunoreactivity. Thus, variations not essentially influencing the immunogenicity of the polypeptide compared to the wild type polypeptide as depicted in SEQ ID NO: 2 or 3 are considered to fall within the scope of the invention. Those variations in the amino acid sequence of a certain structural subunit X or Y according to the invention that still provide a polypeptide capable of inducing an immune response against infection with H. felis or at least against the clinical manifestations of the infection are considered as "not essentially influencing the immunogenicity".

30

35

40

45

50

25

When a polypeptide is used for e.g. vaccination purposes or for raising antibodies, it is however not necessary to use the whole polypeptide. It is also possible to use a fragment of that polypeptide that is capable, as such or coupled to a carrier such as e.g. KLH, of inducing an immune response against that polypeptide, a so-called immunogenic fragment. An "immunogenic fragment" is understood to be a fragment of the full-length polypeptide of the structural subunit X or Y, that still has retained its capability to induce an immune response in the host, i.e. comprises a B- or T-cell epitope. At this moment, a variety of techniques is available to easily identify DNA fragments encoding antigenic fragments (determinants). The method described by Geysen et al (Patent Application WO 84/03564, Patent Application WO 86/06487, US Patent NR. 4,833,092, Proc. Natl Acad. Sci. 81: 3998-4002 (1984), J. Imm. Meth. 102, 259-274 (1987), the so-called PEPSCAN method is an easy to perform, quick and wellestablished method for the detection of epitopes; the immunologically important regions of the polypeptide. The method is used world-wide and as such well-known to man skilled in the art. This (empirical) method is especially suitable for the detection of B-cell epitopes. Also, given the sequence of the gene encoding any protein, computer algorithms are able to designate specific polypeptide fragments as the immunologically important epitopes on the basis of their sequential and/or structural agreement with epitopes that are now known. The determination of these regions is based on a combination of the hydrophilicity criteria according to Hopp and Woods (Proc. Natl. Acad. Sci. 78: 38248-3828 (1981)), and the secondary structure aspects according to

Printed: 10-07-2001

Sequence 3 1 2 3 3 3 3 3 3 3 3 3				gdtdlwaevehdyttygeelkfgagktıregmgqsnspdentldlvitnamiidytgiykadıgikngkihgıgkagnkdmqdgysphmvvgygtea gdtdlwaevehdyttygeelkfgagktıregmgqsnspdentldlvitnamiidytgiykadıgikngkihgıgkagnkdmqdgysphmvvgygtea	qqfptalangvttmfgggtgpvdgtnattitpgkwnlhrmlraaeeysmnvgflgkgnssskkqlvegveagalgfklhedwgttpsaidhclsvade i i i f.s i f.s	namngraihayhiegaggghspdvitmagelnilpssttptipytintvaehldmlmtchhldkriredlqfsqsrirpgslaaedvlhdmgviamts 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	nkkefgklpedgkdndnfrikryiskytinpalthgvseyigsveegkiadlvvwnpaffgvkpkivikggmvvfsemgdsnasvptpgpvyyremfg sa a a r. k. ekg r. r. r. r.
Pequence 2: ureYkuka 1 - Sequence 3: ureYbS4 1 - Sequence 4: ureY301 1 - Sequence 5: ureY301 1 - Sequence 6: B felis 1 - Sequence 6: B felis 1 - Sequence 7: B pylori 1 - Sequence 8: B heilman 1 - Global Protein 1 - Mismatch 2: Open Gap 4 (1) mkmkq-eyvntygptkgdkvrl 1 - I) kisrk sm t r II lo laggamitaggulaptrightytytytytytytytytytytytytytytytytytyty	Homology	99999999999999999999999999999999999999	onserv N	1kfgagktıregmg	pvdgtnattitp	spdvitmageln	kryiskytinpalthgvseyigs
Pequence 2: ureYkuka 1 - Sequence 3: ureYbS4 1 - Sequence 4: ureY301 1 - Sequence 5: ureY301 1 - Sequence 6: B felis 1 - Sequence 6: B felis 1 - Sequence 7: B pylori 1 - Sequence 8: B heilman 1 - Global Protein 1 - Mismatch 2: Open Gap 4 (1) mkmkq-eyvntygptkgdkvrl 1 - I) kisrk sm t r II lo laggamitaggulaptrightytytytytytytytytytytytytytytytytytyty			1;	hdyttygee	ttmfgggtg	hiegagggh:	ykdndnfri) sa sa ca ca kg
Pequence 2: ureYkuka 1 - Sequence 3: ureYbS4 1 - Sequence 4: ureY301 1 - Sequence 5: ureY301 1 - Sequence 6: B felis 1 - Sequence 6: B felis 1 - Sequence 7: B pylori 1 - Sequence 8: B heilman 1 - Global Protein 1 - Mismatch 2: Open Gap 4 (1) mkmkq-eyvntygptkgdkvrl 1 - I) kisrk sm t r II lo laggamitaggulaptrightytytytytytytytytytytytytytytytytytyty	В		xtend	waeve	alangv	raihay ttf tm.tf	Jklped
Pequence 2: ureYkuka 1 - equence 3: ureYbS4 1 - equence 3: ureYbS4 1 - equence 5: ureYbS4 1 - equence 6: B felis 1 - equence 6: B felis 1 - equence 7: B pylori 1 - equence 8: B heilman 1 - equence 9: B heilman 1 - equence	99	υννφυνν οσοσοσο	••	1gdtd	qqfpt	naming:	nkkefg
rs: molecule: cequence 2: cequence 3: cequence 4: cequence 6: cequence 6: cequence 6: cequence 6: cequence 7: cequence 7: cequence 8: cequence 7: cequ	 	1111111	ein Open Gap	-ygptkgdkvr 	jidshthflsp	meagyvddtlr	iprtwqtadkr
molecular construction of the construction of	ureYCS1	ureYkuka ureYDS4 ureY2301 ureY390 B felis B pylori B heilman	Global Prote Mismatch 2;	mkmkq-eyvn:	lagegmiitag	ydvqvcihtdtv	sdsqamgragev
Reference molecul Sequence Seq	:	264696		2222223	120) 120) 120) 120) 121)	240) 240) 240) 240) 241) 241)	3860) 3860) 3861) 360)
Reference mu Sequu UreY300 B felis B beliman ureY2301 ureY3300 B felis B beliman beliman beliman beliman ureY2301 ureY2301 ureY2301 ureY2301 ureY3300 B felis	olecul	ance ance	/pe:				
	Reference m	Nedne Sedne	Alignment ty Parameters:	ureYCS1 ureYkuka ureY2301 ureY230 B felis B pylori B heilman		ureYCS1 ureYkuka ureYDS4 ureY3301 ureY390 B felis B pylori B heilman	ureYCS1 ureYbS4 ureY2301 ureY390 B felis B pylori B heilman

Figure 1c (1)

ALIGNED SEQUENCES



	Номоlоду	1008 998 998 508 528	Global Protein Mismatch 2: Open Gap 4; Extend Gap 1; Conserv N Mismatch 2: Open Gap 4; Extend Gap 1; Conserv N 1) vkltpkeqekfllyyagevarkrkaeglklnqpealayisahimdearrgkktvaqlmeecmhflkkdevmpgvgnmvpdlgveatfpdgtklvtvnwpiepdehfkagevkfgcdkdie 1)	
	(226 aa)	(226 aa) (226 aa) (226 aa) (237 aa) (238 aa) (234 aa)	Extend Gap 1; Conserv N eglkingpealayisahimdearrg r.v.yt.v.l.grvek.d.k.i.yt.v.l.grvek.d.k.i.yt.v.l.e.a.k.i.yt.v.l.ce.a.k.i.yt.v.l.arve.a.a.k.i.yt.v.l.arve.a.a.k.i.yt.v.l.arve.a.a.x.l.d.asfca	
	226	226 226 226 226 237 237 238 234	4, Ex 	
QUENCES	1 H	111111 	Global Protein Mismatch 2; Open Gap 4 wkltpkeqekfllyyagevarkı " 1d 1m , h rl ee m 1d 1m , h l kq m 1d 1m , h l kq " 1d 1m , h l kq	
ALIGNED SEQUENCES	ureXCS1	ureXbS4 ureXDS4 ureX2301 ureX390 A felis A pylori A heilman		
	1e:	8 7 6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	122222222222222222222222222222222222222	
11/15/99	Reference molecule:	Sequence Sequence Sequence Sequence Sequence Sequence Sequence	Alignment type: Parameters: urexCs1 urexCs4 urexCs301 urexC301 urexC301 urexC301 A felis A heilman urexCs1 urexCs1 urexCs1 urexCs1 urexCs300 A felis A felis A heilman urexC301 A heilman A felis A heilman	
			•	

Figure 1



002023333(17.07/20<mark>00</mark>)



7

Chou and Fasman (Advances in Enzymology 47: 45-148 (1987) and US Patent 4,554,101). T-cell epitopes can likewise be predicted from the sequence by computer with the aid of Berzofsky's amphiphilicity criterion (Science 235, 1059-1062 (1987) and US Patent application NTIS US 07/005,885). A condensed overview is found in: Shan Lu on common principles: Tibtech 9: 238-242 (1991), Good et al on Malaria epitopes; Science 235: 1059-1062 (1987), Lu for a review; Vaccine 10: 3-7 (1992), Berzowsky for HIV-epitopes; The FASEB Journal 5:2412-2418 (1991).

- Vaccines against e.g. Helicobacter pylori, which has only one urease, can be made on the basis of this urease, as was described above. In the specific case of Helicobacter felis however a vaccine based upon the known Helicobacter felis structural subunits ureA and B is not capable of providing sufficient protection against Helicobacter felis infection: immunity against structural subunits ureA and B allegedly does not neutralise the urease activity of the newly found heterologous structural subunits UreX and Y.
- Therefore, vaccines for the protection of animals against *Helicobacter felis* infection should at least be directed against the novel urease XY.

 Therefore, one form of still another embodiment of the invention relates to vaccines capable of protecting mammals such as dogs and cats against *Helicobacter felis* infection, that comprise the structural subunit X or Y, preferably X and Y, more preferably X, Y, A and B, or an immunogenic fragment of X and/or Y according to the invention together with a pharmaceutically acceptable carrier.
- Still another embodiment of the present invention relates to the polypeptides according to the invention for use in a vaccine.
 - In still another embodiment, the polypeptide according to the invention is for use in the manufacturing of a vaccine for combating *Helicobacter felis* infections.
- One way of making a vaccine according to the invention is by biochemical purification of the ureaseXY polypeptide or its subunits from a bacterial culture. This can e.g. be done by centrifugation of the bacteria, and the use of gel-filtration columns for separation of the urease polypeptide or its subunits from other components. Further purification may e.g. be done by selective precipitation in ammonium-sulphate, followed by centrifugation, gel electrophoresis and, if desired, separation from the urease AB subunits and dissolving the pellet in a suitable buffer. This is however a time-consuming way of making the vaccine, especially where Helicobacter felis is difficult to grow.
- It is therefore much more convenient to use the expression products of the genes encoding the urease X and Y subunits according to the invention in vaccines. Such vaccines can easily be made by admixing ureaseXY or an UreX or Y subunit or an immunological fragment thereof according to the invention with a pharmaceutically acceptable carrier as described below.
- Furthermore vaccines can comprise live recombinant carriers as described above, capable of expressing ureaseXY, an UreX or UreY subunit or immunogenic fragments thereof according to the invention. Such vaccines, e.g. based upon a Salmonella carrier or a viral carrier infecting the gastric epithelium have the advantage over subunit vaccines that they better mimic the natural way of infection of *Helicobacter felis*.
- Moreover, their self-propagation is an advantage since only low amounts of the recombinant carrier are n cessary for immunisation.



- Vaccines described above all contribute to active vaccination, i.e. the host's immune system is triggered by the UreX and/or Y polypeptide or immunogenic fragments thereof, to make antibodies against these polypeptides.
- Alternatively, such antibodies can be raised in e.g. rabbits or can be obtained from antibody-producing cell lines as described below. Such antibodies can then be administered to the host animal. This method of vaccination, passive vaccination, is the vaccination of choice when an animal is already infected, and there is no time to allow the natural immune response to be triggered. It is also the preferred method for
- vaccinating immune-compromised animals. Administered antibodies against Helicobacter UreX or UreY can in these cases bind directly to the urease excreted by the bacteria. This has the advantage that the urease activity is directly eliminated, thus resulting in acidification of the environment and decreased or stopped Helicobacter growth.
- Therefore, one other form of this embodiment of the invention relates to vaccines comprising antibodies against *Helicobacter felis* urease X polypeptides that have an amino acid sequence that is at least 85 % homologous to SEQ ID NO: 2 or immunogenic fragments of that polypeptide with a length of at least 40 amino acids that are capable of inducing an immune response against ureaseXY or antibodies against
- 20 Helicobacter felis urease Y polypeptides that have an amino acid sequence that is at least 85 % homologous to SEQ ID NO: 3 or immunogenic fragments of that polypeptide with a length of at least 40 amino acids that are capable of inducing an immune response against ureaseXY.
- Vaccines can also be based upon host cells as described above, that comprise ureaseXY, an UreX or UreY subunit or immunogenic fragments thereof according to the invention.
- An alternative and efficient way of vaccination is direct vaccination with DNA encoding the relevant antigen. Direct vaccination with DNA encoding polypeptides has been successful for many different polypeptides. (As reviewed in e.g. Donnelly et al., The Immunologist 2: 20-26 (1993)).
 - This way of vaccination is very attractive for the vaccination of both cats and dogs against *Helicobacter felis* infection.
- Therefore, still other forms of this embodiment of the invention relate to vaccines comprising nucleic acid sequences encoding a polypeptide according to the invention or immunogenic fragments thereof according to the invention, and to vaccines comprising DNA fragments that comprise such nucleic acid sequences.
- Still other forms of this embodiment relate to vaccines comprising recombinant DNA molecules according to the invention.
 - DNA vaccines can easily be administered through intradermal application e.g. using a needle-less injector. This way of administration delivers the DNA directly into the cells of the animal to be vaccinated. Amount of DNA in the microgram range between 1 and 100 µg provide very good results.
- In a further embodiment, the vaccine according to the present invention also comprises antigens from other dog or cat pathogenic organisms and viruses, or genetic information encoding such antig ns. Such organisms and viruses are .g. Feline Infectious Peritonitis virus, Feline Immune deficiency virus, Canine and Feline Parvovirus,
- Distemper virus, Ad novirus, Calicivirus, Bordetella bronchiseptica, Borrelia burgdorferi, Leptospira interrogans, Chlamydia and Bartonella henseli.







Also, the present invention relates to polypeptides according to the invention for use in the manufacturing of a vaccine for combating *Helicobacter felis* infections.

- All vaccines according to the present invention comprise a pharmaceutically acceptable carrier. A pharmaceutically acceptable carrier can be e.g. sterile water or a sterile physiological salt solution. In a more complex form the carrier can e.g. be a buffer.
- Vaccines according to the present invention may in a preferred presentation also contain an adjuvant. Adjuvants in general comprise substances that boost the immune response of the host in a non-specific manner. A number of different adjuvants are known in the art. Examples of adjuvants are Freunds Complete and Incomplete adjuvant, vitamin E, non-ionic block polymers, muramyldipeptides, Quill A^(R), mineral oil e.g. Bayol^(R) or Markol^(R), vegetable oil, and Carbopol^(R) (a homopolymer), or Diluvac^(R) Forte.
- The vaccine may also comprise a so-called "vehicle". A vehicle is a compound to which the polypeptide adheres, without being covalently bound to it. Often used vehicle compounds are e.g. aluminium hydroxide, -phosphate or -oxide, silica, Kaolin, and Bentonite.
- A special form of such a vehicle, in which the antigen is partially embedded in the vehicle, is the so-called ISCOM (EP 109.942, EP 180.564, EP 242.380)
 In addition, the vaccine may comprise one or more suitable surface-active compounds or emulsifiers, e.g. Span or Tween.
- Often, the vaccine is mixed with stabilisers, e.g. to protect degradation-prone polypeptides from being degraded, to enhance the shelf-life of the vaccine, or to improve freeze-drying efficiency. Useful stabilisers are i.a. SPGA (Bovarnik et al; J. Bacteriology 59: 509 (1950)), carbohydrates e.g. sorbitol, mannitol, trehalose, starch, sucrose, dextran or glucose, proteins such as albumin or casein or degradation products thereof, and buffers, such as alkali metal phosphates.
- In addition, the vaccine may be suspended in a physiologically acceptable diluent.

 It goes without saying, that other ways of adjuvating, adding vehicle compounds or diluents, emulsifying or stabilising a polypeptide are also embodied in the present invention.
- Vaccines according to the invention that comprise the UreX or UreY subunit polypeptide can very suitably be administered in amounts ranging between 1 and 100 micrograms, although smaller doses can in principle be used. A dose exceeding 100 micrograms will, although immunologically very suitable, be less attractive for commercial reasons.
- Vaccines based upon live attenuated recombinant carriers, such as the LRC-viruses and bacteria described above can be administered in much lower doses, because they multiply themselves during the infection. Therefore, very suitable amounts would range between 10³ and 10⁹ CFU/PFU for respectively bacteria and viruses.
- Many ways of administration can be applied. Intranasal application is a frequently used way of administrating a vaccine. Oral application is also an attractive way of administration, because the infection is often located in the upper digestive tract. A preferred way of oral administration is the packaging of the vaccine in capsules, known and frequently used in the art, that only disintegrate in the highly acidic environment of the stomach. Also, the vaccine could be mixed with compounds known in the art for

temporarily enhancing the pH of the stomach.

20

25

30

35



10

Systemic application is also suitable, e.g. by intramuscular application of the vaccine. If this route is followed, standard procedures known in the art for systemic application are well-suited.

- Another embodiment of the invention relates to diagnostic tests for the detection of *H. felis* infection. It is known that several *Helicobacter* species such as *H. bizzozeronii*, *H. felis* and *H. salomonis* are capable of infecting both cats and dogs. Of these three, *H. felis* is the species suspected to cause most of the pathology, although it is often outnumbered by *H. bizzozeronii* and *H. salomonis*. Thus, a quick and correct diagnosis of disease, in both cats and dogs, caused by *Helicobacter felis* is important. It has however been very difficult to discriminate between these three species due to the fact that they are so very closely related.
 - Therefore it is another objective of this invention to provide such diagnostic tools suitable for discriminating *H. felis* from other *Helicobacter* species.

On the basis of the novel urease polypeptides and the genes encoding the urease polypeptides, at least three different diagnostic tests, specifically suitable for the discrimination of *H. felis* from other members of the *Helicobacter* family were developed:

1) a diagnostic test based upon the presence or absence of DNA encoding the specific UreX and UreY structural subunits

- 2) a diagnostic test based upon the detection of antibodies against the specific UreX and UreY structural subunits
- 3) a diagnostic test based upon the detection of antigenic material of the specific UreX and UreY structural subunits

A diagnostic test according to 1) is e.g. based upon the reaction of bacterial DNA isolated from the animal to be tested, with specific probes or PCR-primers based upon the sequence of ureX or Y genes. If *H. felis* DNA is present in the animal, this will e.g. specifically bind to ureX or Y specific PCR-primers and will subsequently become amplified in PCR-reaction. The PCR-reaction product can then easily be detected in DNA gel electrophoresis.

The DNA can most easily be isolated from the micro-organisms present in swabs of the upper digestive tract or in the saliva of the animal to be tested. Specific primers can easily be selected from the many regions of the ureX and ureY coding sequences and the non-coding intergenic sequence that differ in sequence from the comparable regions in the ureAB coding sequences. One of the many algorithms suitable for the determination of the level of nucleic acid homology and for comparison of nucleotide sequences in general is known as "Clustal W". It has been described by Thompson et al., in Nucleic Acid Research 22: 4673-4680 (1994). The program can be found at

- several sites on Internet. An more recent alternative for this program is e.g. Align Plus for Windows, available from Scientific and Educational Software, P.O.Box 72045 Durham, NC 27722-2045, USA.
 - As follows from figure 1, a large number of possible PCR-primers can be found that are specific for ureX or ureY. An extremely specific pair of PCR-probes is e.g. formed by the
- 5'-located sequence CATGCACTTTTTGAAAAAAGA (SEQ ID NO: 16) and the 3'-located sequence TATGGTGGTCTTCTCT (SEQ ID NO: 17). Of course many other sequences that are specific for ureX or Y or the intergenic region are suitable. Standard PCR-textbooks give methods for determining the suitability of the probes for selective PCR-reactions with ureX or ureY. PCR-techniques are extensively described in
- 50 (Dieffenbach & Dreksler; PCR primers, a laboratory manual. ISBN 0-87969-447-5 (1995)).

Printed: 10-07-2001







- Another DNA-based test is based upon growth of bacterial material obtained from the swab, followed by classical DNA purification followed by classical hybridisation with radioactively or colour-labelled ureXY-specific DNA-fragments. Given the very low homology between the ureXY-coding regions and the ureAB coding regions of both *H. felis* and other *Helicobacter* species, hybridisation unambiguously indicates the presence or absence of *H. felis*. Both PCR-reactions and hybridisation reactions are well-known in the art and are i.a. described in Maniatis/Sambrook (Sambrook, J. *et al.* Molecular cloning: a laboratory manual. ISBN 0-87969-309-6).
- Selective detection with PCR-primers or with classical hybridisation with ureXY-specific 10 DNA-fragments can be done with fragments that preferably are short, but for practical reasons preferably consist of a stretch of at least 10 contiguous nucleotides of SEQ ID NO: 1. It is clear that for hybridisation experiments a probe needs to be selected that has a higher homology to SEQ ID NO: 1, than to sequences encoding the Helicobacter ureA 15 or ureB subunit. Such a probe can very easily be selected with the help of the Alian Plus for Windows program or the Clustal W program as discussed above. In a comparative hybridisation experiment the DNA to be diagnosed can be tested next to e.g. H. pylori DNA. The probe according to the invention, having a higher homology to SEQ ID NO: 1 than to a gene encoding ureAB, would bind better to H. felis DNA (if present in the 20 sample) than to DNA of other Helicobacter species thus specifically revealing the presence of H. felis DNA in the sample to be tested. The sequences SEQ ID NO: 16 or 17 mentioned above are merely examples of probes very suitable for labelling and
- 25 Thus, one embodiment of the invention relates to a diagnostic test for the detection of DNA encoding the specific *Helicobacter* UreX and UreY subunit polypeptides. Such a test comprises a nucleic acid sequence according to the invention or a fragment thereof that is specific for the DNA encoding UreX and UreY or the intergenic region between UreX and UreY. A fragment that is specific for that DNA is a fragment that binds better to the DNA encoding UreX and UreY or the intergenic region between UreX and UreY than to the DNA encoding UreA and UreB or the intergenic region between UreA and UreB.

subsequent use in the H. felis-specific hybridisation assays as described.

- Methods for the detection of *Helicobacter felis* DNA comprise hybridisation of the DNA to be tested with UreX or Y DNA, or PCR-reaction of the DNA to be tested with UreX or Y DNA specific probes.
- A diagnostic test according to 2) for the detection of *Helicobacter felis* antibodies in sera can be e.g. a simple sandwich-ELISA-test in which purified UreX or UreY subunit polypeptides or antigenic fragments thereof according to the invention are coated to the wall of the wells of an ELISA-plate. A method for the detection of such antibodies is e.g. Incubation of purified UreX or Y polypeptide with serum from mammals to be tested, followed by e.g. incubation with a labelled antibody against the relevant mammalian antibody. A colour reaction can then reveal the presence or absence of antibodies against *Helicobacter felis* urease XY. Depending on the labelled antibodies used, the selectivity of this system can be improved by pre-incubation of the serum to be tested with urease AB followed by spinning down the precipitate, in order to avoid non-XY-
 - If antigenic fragments of the UreX or UreY structural subunits according to the invention are used for coating, this pre-incubation step can be skipped.

specific reactions.



00202565.8(17-07-2000)



12

Another example of a diagnostic test system is e.g. the incubation of a Western blot comprising UreX or UreY polypeptide or an antigenic fragment thereof according to the invention, with serum of mammals to be tested, followed by analysis of the blot. The purified UreX and UreY structural subunits or antigenic fragments thereof according to the invention, suitable for the coating of ELISA plates or for Western blotting can easily be obtained by expression of the ureX and ureY gene as was described by Ferrero for ureA and B (Ferrero et al., Molec. Microbiol. 9, 323-333 (1993)).

Also, the invention relates to methods for the detection in serum of antibodies against

Helicobacter felis antibodies in which the method comprises the incubation of serum with

UreX or UreY polypeptide or an antigenic fragment thereof according to the invention.


A diagnostic test according to 3) based upon the detection of antigenic material of the specific UreX and UreY structural subunits of Helicobacter felis antigens and therefore suitable for the detection of Helicobacter felis infection can e.g. also be a standard 15 ELISA test. In one example of such a test the walls of the wells of an ELISA plate are coated with antibodies directed against the specific UreX and UreY structural subunits of Helicobacter felis. The antigenic material to be tested can if necessary be pre-incubated with antibodies against UreA and B. This will leave the UreX and Y specific epitopes 20 uncovered and therefore the pre-incubated Helicobacter species will bind to the ELISA plate only if it comprises UreX or Y, i.e. if it specifically is Helicobacter felis. The use of monoclonal antibodies specific for UreX or Y and not reacting with UreA or B are the preferred antibodies in such tests, because they make the pre-incubation step superfluous. Such monoclonal antibodies can easily be obtained by immunising inbred 25 mice with immunising fragments of UreX or Y according to the invention, by techniques also known in the art (See below: Kohler and Milstein).

The polypeptides or immunogenic fragments thereof according to the invention expressed as characterised above can be used to produce antibodies, which may be polyclonal, monospecific or monoclonal (or derivatives thereof). If polyclonal antibodies are desired, techniques for producing and processing polyclonal sera are well-known in the art (e.g. Mayer and Walter, eds. *Immunochemical Methods in Cell and Molecular Biology*, Academic Press, London, 1987).

Monoclonal antibodies, reactive against the polypeptide according to the invention (or variants or fragments thereof) according to the present invention, can be prepared by immunising inbred mice by techniques also known in the art (Kohler and Milstein, *Nature*, 256, 495-497, 1975).

Finally, the invention relates to methods for the detection of antigenic material from Helicobacter felis in which the method comprises the incubation of serum, tissue of body fluids with antibodies against UreX or UreY polypeptide or an antigenic fragment thereof according to the invention.







Example 1

The ureX and ureY genes of Helicobacter felis strain CS1: cloning and expression in Escherichia coli.

The *ureX* and *ureY* genes of *H. felis* strain CS1 were cloned as an operon into an *E. coli* T7 expression vector, pET3a, as follows:

- For proper expression of the UreX and Y proteins in pET3a (Novagen, 601 Science 10 Drive, Madison WI, USA) the genes were cloned as a Ndel-BamHI DNA fragment into the Ndel-BamHI sites of this vector. The ureaseXY operon contains an internal Ndel site that was mutated by overlap-extension PCR of 2 PCR fragments. For that purpose two PCR fragments (the 5' and the 3' products) were amplified using chromosomal DNA of H. felis CS1 as the template. The 5' PCR product contained the complete ureX gene and the first part of the ureY gene. The forward primer contained a Ndel restriction site and 15 the start codon of ureX (GGAGTAACATATGAAACTCACA CCCAAAGAGC) (SEQ ID and the reverse primer contains a point mutation (CACACCC ACGACCATGTGAGGGCTTAC) (SEQ ID NO: 19). The second, 3' PCR product consisted of the 3' end of the ureY gene. This forward primer is complementary to the reverse primer of the first PCR product and also contained the same point mutation 20 (GTAAGCC CTCACATGGTCGTGGGTGTG) (SEQ ID NO: 20), and the reverse primer contained a BamHI restriction site just downstream of the stopcodon of the ureY gene (CGAATT CGGATCCTAGAAGAAAGTGTAGCGCTGG) (SEQ ID NO: 21). The mutation in the complementary primers is made to delete the internal Ndel site in ureY, it replaces the CATATG (His-Met) by CACATG (His-Met).
- the CATATG (His-Met) by CACATG (His-Met).

 After amplification of both PCR products, the complete operon was obtained by overlap-extension-PCR with the forward primer of the *ureX* and the reverse primer of the *ureY* using both PCR products as templates. The resulting PCR product was cloned into PCR-bluntII-TOPO (Invitrogen, P.O.Box 2312, 9704 CH Groningen, The Netherlands) and transformed into *E. coli* TOP10F' cells (Invitrogen). Positive clones were isolated and the *ureaseXY* genes were sub-cloned into pET3a with *NdeI-BamHI*. The obtained plasmid was called pUreXY-1 and was transformed into the expression strain HMS174(DE3)/pLysS (Novagen).
- The ureX and ureY genes of pUreXY-1 were expressed in HMS174(DE3)/pLysS as follows: an overnight culture was diluted 1/100 into TB Amp¹⁰⁰ Cam²⁵; this culture was incubated for 3 h at 37°C at 200 rpm; the culture was induced by adding 1 mM of IPTG and incubated for another 3 h at 37°C at 200 rpm. The induction was done twice, once in a small scale and once in a large scale.
- The induced samples were analysed on a SDS-PAGE gel (fig. 2). As can be clearly seen from lane 9, expression of UreX and UreY, when induced provides the two structural subunits as polypeptide bands with a molecular weight of 25 kDa for the UreX subunit and 62 kDa for the UreY subunit.



00202565,8(17-07-2000)



14

Legend to the figures

- Figure 1a: Comparison of the nucleic acid sequence encoding UreX and Y, including a short non-coding region bridging the two coding sequences, from *Helicobacter felis* species CS1, Kukka, Ds4, 2301 and 390 with the nucleic acid sequence encoding UreA and B, including a short non-coding region bridging the two coding sequences, from *Helicobacter felis*, *pylori* and *heilmannii*
- Figure 1b: Comparison of the amino acid sequence of UreX from Helicobacter felis species CS1, Kukka, Ds4, 2301 and 390 with the amino acid sequence encoding UreA from Helicobacter felis, pylori and heilmannii
- Figure 1c: Comparison of the amino acid sequence of UreY from *Helicobacter felis* species CS1, Kukka, Ds4, 2301 and 390 with the amino acid sequence encoding UreB from *Helicobacter felis*, *pylori* and *heilmannii*

Figure 2: Polyacrylamide gel of the expression products UreX and UreY

20	Lane 7 Lane 8 Lane 9 Lane 10	 : Biorad broad range marker : Complete cell culture before induction (small scale culture) : Complete cell culture after induction (small scale culture) : Complete cell culture after induction (large scale culture)
25	Lane 11	: Supernatant after induction (large scale culture). : Biorad pre-stained marker



against ureaseXY.

- 12) Polypeptide according to claims 8-11 fur use in a vaccine
- 13) Polypeptide according to claims 8-11 for use in the manufacturing of a vaccine for combating Helicobacter felis infections.
- 14) Vaccine for combating *Helicobacter felis* infections, characterised in that it comprises a nucleic acid sequence according to claims 1-3, a DNA fragment according to claim 4, a recombinant DNA molecule according to claim 5, a live recombinant carrier according to claim 6, a host cell according to claim 7 or a polypeptide according to claims 8-11, and a pharmaceutically acceptable carrier.
- 15) Vaccine according to claim 14, characterised in that it comprises an adjuvant.
- 16) Vaccine according to claim 14 or 15, characterised in that it comprises an additional antigen derived from a virus or micro-organism pathogenic to mammals or genetic information encoding said antigen.
- 17) Vaccine according to claim 16, characterised in that said virus or micro-organism pathogenic to mammals is selected from the group of Feline Infectious Peritonitis virus, Feline Immune deficiency virus, Canine and Feline Parvovirus, Distemper virus, Adenovirus, Calicivirus, Bordetella bronchiseptica, Borrelia burgdorferi, Leptospira interrogans, Chlamydia and Bartonella henseli.
- 18) Vaccine for combating *Helicobacter felis* infections, characterised in that it comprises antibodies against a polypeptide according to claims 8-11.
- 19) Method for the preparation of a vaccine according to claims 14-17, said method comprising the admixing of a polypeptide according to claims 8-11 and a pharmaceutically acceptable carrier.
- 20) Diagnostic test for the detection of Helicobacter felis specific DNA characterised in that the test comprises a nucleic acid sequence according to claims 1-3, or a fragment thereof.
- 21) Diagnostic test for the detection of antibodies against *Helicobacter felis*, characterised in that said test comprises a polypeptide or a fragment thereof as described in claims 8-11.
- 22) Diagnostic test for the detection of antigenic material of Helicobacter felis, characterised in that said test comprises antibodies against a polypeptide or a fragment thereof as described in claims 8-11.



00202565.8(17-07-2000)



EPO - DG 1

17. 07. 2000

Claims

- 1) Nucleic acid sequence encoding two subunit polypeptides of a urease complex such as expressed by *Helicobacter felis*, said nucleic acid sequence having at least 85 % homology with SEQ ID NO: 1, or a part thereof encoding at least an immunogenic fragment of one of said subunits, said part having a length of at least 40, preferably 45, more preferably 50 nucleotides.
- 2) Nucleic acid sequence according to claim 1, characterised in that it encodes the urease X subunit polypeptide or the urease Y subunit polypeptide.
- 3) Nucleic acid sequence according to claim 1 or 2, characterised in that the sequence has at least 90 %, preferably 94 %, more preferably 97 % homology therewith.
- 4) DNA fragment comprising a nucleic acid sequence according to claims 1-3
- 5) Recombinant DNA molecule comprising a nucleic acid sequence according to claims 1-3 or a DNA fragment according to claim 4, under the control of a functionally linked promoter.
- 6) Live recombinant carrier comprising a recombinant DNA molecule according to claim 5
- 7) Host cell comprising a nucleic acid sequence according to claims 1-3, a DNA fragment according to claim 4, a recombinant DNA molecule according to claim 5 or a live recombinant carrier according to claim 6.
- 8) Helicobacter felis urease X subunit polypeptide, said polypeptide having an amino acid sequence that is at least 85 % homologous to SEQ ID NO: 2 or an immunogenic fragment of said polypeptide with a length of at least 40, preferably 45, more preferably 50 amino acids said immunogenic fragment being capable of inducing an immune response against ureaseXY.
- 9) Polypeptide according to claim 8, having a sequence homology of at least 85 %, preferably 90 %, more preferably 95 % homology to SEQ ID NO: 2, or an immunogenic fragment of said polypeptide capable of inducing an immune response against ureaseXY.
- 10) Helicobacter felis urease Y subunit polypeptide, said polypeptide having an amino acid sequence that is at least 85 % homologous to SEQ ID NO: 3 or an immunogenic fragment of said polypeptide with a length of at least 40, preferably 45, more preferably 50 amino acids said immunogenic fragment being capable of inducing an immune response against ureaseXY.
- 11) Polypeptide according to claim 10, having a sequence homology of at least 85 %, preferably 90 %, more preferably 95 % homology to SEQ ID NO: 3, or an immunogenic fragment of said polypeptide capable of inducing an immune response



00202565.8(17-07-2000)



17

EPO - DG 1

17. 07. 2000

Abstract

The present invention relates to novel *Helicobacter felis* urease subunit polypeptides and to nucleic acid sequences encoding these subunit polypeptides, to DNA fragments and recombinant DNA molecules comprising the nucleic acid sequences encoding these subunit polypeptides, to live recombinant carriers and to host cells comprising nucleic acid sequences encoding these subunit polypeptides. Also, the invention relates to the subunit polypeptides for use in vaccines and for use in the manufacturing thereof, to vaccines comprising said subunit polypeptides and to methods for the preparation of such vaccines. Furthermore, the invention relates to diagnostic methods for the detection of *Helicobacter felis* specific nucleic acid sequences, *Helicobacter felis* antigenic material and to antibodies against *Helicobacter felis*.

		·		•
				P
			,	
			·	į
·				
	•			
	,			
•				
				·
·				
				•

EPO	-	DG	1

17. 07. 2000

948 948 858 678 678

2398 bps) 2398 bps) 2398 bps) 2179 bps) 2433 bps) 2430 bps) 2426 bps)

2445 2399 2398 2181 2475 5088 2636

48 2 3 3 2659 211

Kukkaseq Ds4seq 2301SEQ 390Seq H. felis comp H. pylori com H. heilmannii

26.4.0.6.8

Sequence Sequence Sequence Sequence Sequence Sequence Global DNA Mismatch 1; Open Gap 3; Extend Gap

Alignment type: Parameters:

Homolog₃

2398 bps)

206 gtgaaactcacacacaaagagcaaagttcttgttatattatgcgggcgaaagtggctagaaagcgcaagcagtcaaccaaaccaaaccaaaccattgcttacattagt	326 gcccatattatggacgagagagagagagagagatgacattattgaaaaaagatgaaatgaatg	446) ctaggtgtagaagccactttcctgatggtacgaaacttgtaacttggaaccagatgagcacttcaaagcgggggaagttgatgttggttagataaagacatcgag 288) t.g.c.gt.c.c.c.c.c.c.c.c.c.c.c.c.c.c.	566 ctcaatgcaggcaaagaactgaagttactaatgaagggcctaaatccttgcatgtgggtagccatttccacttctttgaagctaacaaggcactaaaattcgatcgtgaa 408 ct.g
Cslseq Kukkaseq Dslseq 2301SEQ 390seq H. fells com H. pylori co H. hellmanni	Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co H. heilmanni	Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felia com H. pylori co H. heilmanni	Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co

Figure 1a (

ALIGNED SEQUENCES

Cslseq

Reference molecule:

686) aaagcctatggcaaacgcctctggcaacacgctacgcattggggcaggacaaacccgcaaagtgcagttgattcctcttgggtggcagtaaaaagtgattggcatgaac 528) 482) 482) 483) 481) 483) 483) 483) 484) 520) 485) 487) 487) 520 53136)	806) gggcttgtgaataacatcgcggatga	878)atcaagtaaggagactcccatgaaaatgaaaaaacaagaatatgtaaatacctacggacccaccaaaggcgataaaagtgcgcttaggaga 720) 674) 675) 677) 677) 677) 677) 677) 677) 677) 677) 677) 677) 677) 677) 677) 677) 677)	968) taccgatctttgggcagaagtagaacatgactataccacctatggcgaagaacttagatttggcggggtaaaactatccgtgaggggtatggggcagaatagccctgatgaaaacac 810)	1088) cctagatttagtcatcactaacgcgatgattacaaagccgacattgggattaaaaacggcaaaatccatggcattggcaaggcaaggaaacaaggacatgca 930) t. 884) c. 885) t. 885) t. 885) t. 885) t. 886) t. 887 c. 887 c. 887 c. 888 c. 887 c. 888 c. 888 c. 888 c. 889 c. 889 c. 889 c. 889 c. 889 c. 889 c. 880 c. 8	1208) agatggcgtaagccctcatatggtcgtgggtgtggggcacagaagcactagcaggggaaggtatgattattaccgctgggggaatcgattcaccacactcctttctccacaacaatt 1050) 1004) 1005) 1005) 1005) 1008) 10
Csiseq Kukkaseq Ds4seq 2301SEQ 390seq H. fells com (H. hellmanni (Cslseq Kukkaseq Ds4seq (2301SEQ 390seq H. felis com H. hellmanni (Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. hellmanni (Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. hellmanni (Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co (Cslseq Kukkaseq Ds4seq 2301SEQ 390seq H. felis com H. pylori co (

Figure 1a (

480) hhgkakfdtsitfvskvayengvkeklglerqvlpvkncrnitkkdfkfndktakitvdpktfevfvdgklctskptsqvplagrytff 480) 480) 480) 480) 480) 480) 480) 480)
480) hhq 480) 480) 481)
ureYCS1 ureYbka ureYbS4 ureY3301 ureY390 B felis

Figure 1c (2)




Figure .





SEQUENCE LISTING

EPO-DG 1
17. 07. 2000

<110> AKZO Nobel N.V.

<120> Helicobacter vaccine

<130> Degoedesequenties

<140>

<141>

<160> 15

<170> PatentIn Ver. 2.1

<210> 1

<211> 2405

<212> DNA

<213> Helicobacter felis

<220>

<221> CDS

<222> (1)..(681)

<220>

<221> CDS

<222> (692)..(2398)

<400> 1

gtg aaa ctc aca ccc aaa gag caa qaa aag ttc ttg tta tat tat gcg 48 Val Lys Leu Thr Pro Lys Glu Glu Lys Phe Leu Leu Tyr Tyr Ala 10

ggc gaa gtg gct aga aag cgc aaa gca gag ggc tta aag ctc aat caa 96 Gly Glu Val Ala Arg Lys Arg Lys Ala Glu Gly Leu Lys Leu Asn Gln 20 25

ccc gaa gcc att gcc tac att agt gcc cat att atg gac gag gcg cgc 144 Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala Arq 35

cgt ggc aaa aaa acc gtt gct gaa ctt atg gaa gaa tgt atg cac ttt Arg Gly Lys Lys Thr Val Ala Glu Leu Met Glu Glu Cys Met His Phe 50 55 60

ttg aaa aaa gat gag gtg atg ccc ggt gtg ggg aat atg gtc cct gat Leu Lys Lys Asp Glu Val Met Pro Gly Val Gly Asn Met Val Pro Asp 65 70 75





ttg	ggc	: gta	gaa	gcc	act	ttc	CCC	gat	ggc	acc	aaa	ctc	gta	acc	gtg	288
Leu	Gly	v Val	Glu	Ala	Thr	Phe	Pro	Asp	Gly	Thr	Lys	Leu	Val	Thr	Val	
				85					90		_			95		
aat	taa	CCC	att	αaa	cct	gat	gaa	cac	+++	222	acc	aat	~~~	at a	222	226
												-				336
ASII	ırp	PIO		GIU	PLO	Asp	GLU		Pne	тÀг	ATa	GIA		val	ьуs	
			100					105					110			
ttt	ggc	tgt	gat	aaa	gac	att	gag	ctc	aac	gcg	ggt	aag	gaa	gtt	acc	384
Phe	Gly	Cys	Asp	Lys	Asp	Ile	Glu	Leu	Asn	Ala	Gly	Lys	Glu	Val	Thr	
		115					120					125				
gag	ctt	qaa	gtt	acc	aac	gaa	gga	cct	aaa	tcc	tta	cat.	ata	aat	agc	432
						Glu										102
	130					135	1		_,_		140		·uı	OT A	Der	
	100					133					140					
+	++-		++-													
						acc									_	480
	Pne	HIS	Pne	Pne		Thr	Asn	Lys	Ala			Phe	Asp	Arg	Glu	
145					150					155	•				160	
aaa	gcc	tat	ggc	aaa	cgc	cta	gat	att	ccc	tct	ggc	aac	acg	cta	cgc	528
Lys	Ala	Tyr	Gly	Lys	Arg	Leu	Asp	Ile	Pro	Ser	Gly	Asn	Thr	Leu	Arg	
				165					170					175		
att	ggg	gca	gga	caa	acc	cgt	aaa	ata	caq	tta	atc	cct	ctt	aac	aat.	576
						Arg										0.0
	3		180			5		185	02		110		190	O.L.y	OLY	
								100					100			
- ~+	222		~+~	~++	~~~	- + 		~ ~ ~							_	60.4
						atg									-	624
ser	гуѕ		vaı	шe	GIÀ	Met		GTÅ	Leu	vaı	Asn		lie	Ala	Asp	
		195					200					205				
			•													
						gca										672
Glu	Arg	His	Lys	His	Lys	Ala	Leu	Asp	Lys	Ala	Lys	Ser	His	Gly	Phe	
	210					215					220					
atc	aag	taa	ggag	racto	cc a	itg a	aa a	itg a	aa a	aa c	aa g	ag t	at g	ta a	ac	721
Ile	Lys				M	et L	ys N	1et I	ys I	ys G	ln G	lu T	'yr V	al P	sn	
225						•		230	_	-			35			
							-									
acc	tac	gga	ccc	acc	aca	ggc	gat	aaa	ata	cac	tta	aas	αat	acc	αat	769
						Gly							-		-	109
Y TI T	TÄT	91 y 240	£ 1.0	1117	TIIT			пЛэ	val	ALY	ъeu	_	ush	THE	АЅР	
		240					245					250				
						cat							-	_		817
Leu		Ala	Glu	Val	Glu	His	Asp	Tyr	Thr	Thr	_	Gly	G1 ų	Glu	Leu	
	255					260					265					



Printed:10-07-2001

00202333 3(17 07 2000)



	ttt Phe						_			-		-	-		865
	cca Pro												_		913
	gac Asp			_			_	_							961
	atc Ile					-									1009
-	agc Ser 335			_	-	 				_	-		_		1057
	ggt Gly									-					1105
	tct Ser														1153
	ttt Phe														1201
	act Thr	_				-		-	-	_	-	-	-	_	1249
	tat Tyr 415		_				_					_		_	1297
	aaa Lys			-		_	-								1345
	cat His							_							1393







			tac Tyr						_	_	1441
			tat Tyr				_	Met			1489
			tac Tyr								1537
			atg Met 515								1585
			ccc Pro								1633
			tgc Cys					_		_	1681
			agc Ser								1729
			att Ile								1777
			gct Ala 595			-					1825
			aaa Lys				-	-		-	1873
	Asp		cgc Arg	Lys			Lys				1921
Pro			cat His			Ile					1969





				gct ttc ttt ggt : Ala Phe Phe Gly	2017
•				gtg ttc tct gaa : Val Phe Ser Glu 685	2065
				ccg gtt tat tac 2 Pro Val Tyr Tyr 700	2113
Arg Glu Met E				gac acc agc atc : Asp Thr Ser Ile 715	2161
_	_		Asn Gly Val	aaa gaa aaa cta 2 Lys Glu Lys Leu 730	2209
			_	cgc aac atc act 2 Arg Asn Ile Thr	2257
-		_		atc act gtc gat 2 Ile Thr Val Asp 765	2305
				tgc acc tct aaa 2 Cys Thr Ser Lys 780	2353
ccc gcc tct g Pro Ala Ser G		-	-		2398
gcacaat					2405
<210> 2 <211> 226 <212> PRT <213> Helicoh	bacter feli:	S			
<400> 2 Val Lys Leu 1	Thr Pro Lys 5	Glu Gln Glu	Lys Phe Leu	Leu Tyr Tyr Ala 15	



Gly	Glu	Val	Ala	Arg	Lys	Arg	Lys	Ala	Glu	Gly	Leu	Lys	Leu	Asn	Gln
			20					25					30		

Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala Arg 35 40 45

Arg Gly Lys Lys Thr Val Ala Glu Leu Met Glu Glu Cys Met His Phe 50 55 60

Leu Lys Lys Asp Glu Val Met Pro Gly Val Gly Asn Met Val Pro Asp 65 70 75 80

Leu Gly Val Glu Ala Thr Phe Pro Asp Gly Thr Lys Leu Val Thr Val 85 90 95

Asn Trp Pro Ile Glu Pro Asp Glu His Phe Lys Ala Gly Glu Val Lys
100 105 110

Phe Gly Cys Asp Lys Asp Ile Glu Leu Asn Ala Gly Lys Glu Val Thr 115 120 125

Glu Leu Glu Val Thr Asn Glu Gly Pro Lys Ser Leu His Val Gly Ser 130 135 140

His Phe His Phe Phe Glu Thr Asn Lys Ala Leu Lys Phe Asp Arg Glu 145 150 155 160

Lys Ala Tyr Gly Lys Arg Leu Asp Ile Pro Ser Gly Asn Thr Leu Arg 165 170 175

Ile Gly Ala Gly Gln Thr Arg Lys Val Gln Leu Ile Pro Leu Gly Gly
180 185 190

Ser Lys Lys Val Ile Gly Met Asn Gly Leu Val Asn Asn Ile Ala Asp 195 200 205

Glu Arg His Lys His Lys Ala Leu Asp Lys Ala Lys Ser His Gly Phe 210 215 220

Ile Lys 225

<210> 3

<211> 568

<212> PRT

<213> Helicobacter felis



<4	ດດ	1>	3

- Met Lys Met Lys Lys Gln Glu Tyr Val Asn Thr Tyr Gly Pro Thr Thr

 1 5 10 15
- Gly Asp Lys Val Arg Leu Gly Asp Thr Asp Leu Trp Ala Glu Val Glu
 20 25 30
- His Asp Tyr Thr Thr Tyr Gly Glu Glu Leu Lys Phe Gly Ala Gly Lys
 35 40 45
- Thr Ile Arg Glu Gly Met Gly Gln Ser Asn Ser Pro Asp Glu Asn Thr 50 55 60
- Leu Asp Leu Val Ile Thr Asn Ala Met Ile Ile Asp Tyr Thr Gly Ile
 65 70 75 80
- Tyr Lys Ala Asp Ile Gly Ile Lys Asn Gly Lys Ile His Gly Ile Gly 85 90 95
- Lys Ala Gly Asn Lys Asp Met Gln Asp Gly Val Ser Pro His Met Val 100 105 110
- Val Gly Val Gly Thr Glu Ala Leu Ala Gly Glu Gly Met Ile Ile Thr 115 120 125
- Ala Gly Gly Ile Asp Ser His Thr His Phe Leu Ser Pro Gln Gln Phe 130 135 140
- Pro Thr Ala Leu Ala Asn Gly Val Thr Thr Met Phe Gly Gly Gly Thr 145 150 155 160
- Gly Pro Val Asp Gly Thr Asn Ala Thr Thr Ile Thr Pro Gly Lys Trp
 165 170 175
- Asn Leu His Arg Met Leu Arg Ala Ala Glu Glu Tyr Ser Met Asn Val 180 185 190
- Gly Phe Leu Gly Lys Gly Asn Ser Ser Ser Lys Lys Gln Leu Val Glu 195 200 205
- Gln Val Glu Ala Gly Ala Ile Gly Phe Lys Leu His Glu Asp Trp Gly 210 215 220
- Thr Thr Pro Ser Ala Ile Asp His Cys Leu Ser Val Ala Asp Glu Tyr 225 230 235 240
- Asp Val Gln Val Cys Ile His Thr Asp Thr Val Asn Glu Ala Gly Tyr

_	•	_
2	4	.5

- Val Asp Asp Thr Leu Asn Ala Met Asn Gly Arg Ala Ile His Ala Tyr 260 265 270
- His Ile Glu Gly Ala Gly Gly Gly His Ser Pro Asp Val Ile Thr Met 275 280 285
- Ala Gly Glu Leu Asn Ile Leu Pro Ser Ser Thr Thr Pro Thr Ile Pro 290 295 300
- Tyr Thr Ile Asn Thr Val Ala Glu His Leu Asp Met Leu Met Thr Cys 305 310 315 320
- His His Leu Asp Lys Arg Ile Arg Glu Asp Leu Gln Phe Ser Gln Ser 325 330 335
- Arg Ile Arg Pro Gly Ser Ile Ala Ala Glu Asp Val Leu His Asp Ile 340 345 350
- Gly Val Ile Ala Met Thr Ser Ser Asp Ser Gln Ala Met Gly Arg Ala 355 360 365
- Gly Glu Val Ile Pro Arg Thr Trp Gln Thr Ala Asp Lys Asn Lys Lys 370 375 380
- Glu Phe Gly Lys Leu Pro Glu Asp Gly Ala Asp Asn Asp Asn Phe Arg 385 390 395 400
- Ile Lys Arg Tyr Ile Ser Lys Tyr Thr Ile Asn Pro Ala Leu Thr His
 405 410 415
- Gly Val Ser Glu Tyr Ile Gly Ser Val Glu Glu Gly Lys Ile Ala Asp 420 425 430
- Leu Val Val Trp Asn Pro Ala Phe Phe Gly Val Lys Pro Lys Ile Val 435 440 445
- Ile Lys Gly Gly Met Val Val Phe Ser Glu Met Gly Asp Ser Asn Ala 450 455 460
- Ser Val Pro Thr Pro Gln Pro Val Tyr Tyr Arg Glu Met Phe Gly His 465 470 475 480
- His Gly Lys Ala Lys Phe Asp Thr Ser Ile Thr Phe Val Ser Lys Val
 485 490 495
- Ala Tyr Glu Asn Gly Val Lys Glu Lys Leu Gly Leu Glu Arg Lys Val

505

510

Leu Pro Val Lys Asn Cys Arg Asn Ile Thr Lys Lys Asp Phe Lys Phe 515 520 525

Asn Asn Lys Thr Ala His Ile Thr Val Asp Pro Lys Thr Phe Glu Val 530 535 540

Phe Val Asp Gly Lys Leu Cys Thr Ser Lys Pro Ala Ser Glu Val Pro 545 550 555 560

Leu Ala Gln Arg Tyr Thr Phe Phe 565

<210> 4

<211> 2183

<212> DNA

<213> Helicobacter felis

<220>

<221> CDS

<222> (3)..(683)

<220>

<221> CDS

<222> (694)..(2181)

<400> 4

tc gtg aaa ctc aca ccc aaa gag caa gaa aag ttc ttg tta tat tat 47
Val Lys Leu Thr Pro Lys Glu Gln Glu Lys Phe Leu Leu Tyr Tyr
1 5 10 15

gcg ggc gaa gtg gct aga aag cgc aaa gca gag ggc tta aag ctc aat 95
Ala Gly Glu Val Ala Arg Lys Arg Lys Ala Glu Gly Leu Lys Leu Asn
20 25 30

caa ccc gaa gcc att gcc tac att agt gcc cat att atg gac gag gcg 143 Gln Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala 35 40 45

cgc cgt ggc aaa aaa acc gtt gct gaa ctt atg gaa gaa tgt atg cac 191 Arg Arg Gly Lys Lys Thr Val Ala Glu Leu Met Glu Glu Cys Met His 50 55 60

ttt ttg aaa aaa gat gag gtg atg ccc ggt gtg ggg aat atg gtc cct 239 Phe Leu Lys Lys Asp Glu Val Met Pro Gly Val Gly Asn Met Val Pro

		65					70					75					
As		_	-		-								aaa Lys		_		287
_	-					-		_	-				gcc Ala		-	, ,	335
				-	_		-			Leu			ggt Gly	_	-	-	383
				_	_			_					ttg Leu 140				431
-	r						_			-	-	_	aaa Lys		-		479
-	u		-				-		-				Gly		_		527
-				-				_			-		atc Ile				575
,		-						_					aat Asn				623
-		_	_										aaa Lys 220				671
	e	atc Ile 225	_	taa	ggag	gacto		-	ys I	_			caa q Gln (3lu 7			720
									-			-	tta Leu		_		768
ga	t	ctt	tgg	gca	gaa	gta	gaa	cat	gac	tat	acc	act	tat	ggc	gaa	gag	816

Asp Leu Trp Ala Glu Val Glu His Asp Tyr Thr Thr Tyr Gly Glu Glu

	aaa													_	-	864
Leu	Lys 270	Phe	Gly	Ala	Gly		Thr	Ile	Arg	Glu	-	Met	Gly	Gln	Ser	
•	270					275					280					
aat	agc	cca	gat	gaa	aac	acc	tta	σat	tta	ata	atc	acc	aac	aca	ato	912
	Ser														_	712
285					290					295					300	
										•						
	atc															960
He	Ile	Asp	Tyr	Thr 305	Gly	Ile	Tyr	Lys			Ile	Gly	Ile		Asn	
				303					310					315		
ggc	aaa	atc	cat	ggt	att	ggc	aag	gcg	ggg	aac	aaa	gac	atg	caa	gat	1008
	Lys															
			320					325					330			
													•			
	gta Val															1056
01,	vul	335			1100	vai	340	GIY	vai	GIĀ	1111	345	Ala	ьеи	AIA	
	gaa															1104
Gly	Glu	Gly	Met	Ile	Ile		Ala	Gly	Gly	Ile	Asp	Ser	His	Thr	His	
	350					355					360					
ttc	ctc	tct	ccc	caa	caa	ttc	cct	acc	act	cta	acc	aat	aat	att		1152
	Leu													-		1132
365					370					375			•		380	
	atg													-		1200
Thr	Met	Pne	GIA	385	GLY	Thr	GLY	Pro	Va⊥ 390	Asp	GTA	Thr	Asn	Ala 395	Thr	
				505					390					393		
acc	atc	act	ccg	ggc	aaa	tgg	aac	ttg	cac	cgc	atg	ttg	cgc	gca	gct	1248
Thr	Ile	Thr	Pro	Gly	Lys	Trp	Asn	Leu	His	Arg	Met	Leu	Arg	Ala	Ala	
			400					405					410			
~ > >	~~~	tat	tat	ata	22t	at a	~~~		++ ~	~~~		~~~			.	1200
	gag Glu															1296
		415			- 		420			0_1	-1-	425		002	JU2	
	•															
	aaa															1344
Ser	Lys	Lys	Gln	Leu	Val		Gln	Val	Glu	Ala	_	Ala	Ile	Gly	Phe	
	430					435					440					
aaa.	ttg	cat	gaa	gac	tgg	ggc	aca	aca	сса	agt	aca	atc	gat	cac	tac	1392
	Leu									-			-		-	

445	•			450			455	ı				460	
		gtg Val		Glu			Val					Asp	1440
		aat Asn									_		1488
		gcc Ala 495											1536
		gat Asp											1584
		acc Thr								-	_		1632
		atg Met					-	_	_		_		1680
		caa Gln									_	-	1728
		gtg Val 575								_		-	1776
		gca Ala							_				1824
		gat Asp											1872
gca Ala		aac Asn											1920
att Ile		ccc Pro											1968



640 645 650

gaa gag ggc aag atc gcc gac ttg gtg gtg tgg aat cct gcc ttt ttt 2016 Glu Glu Gly Lys Ile Ala Asp Leu Val Val Trp Asn Pro Ala Phe Phe 655 660 665

ggc gtg aaa cct aag att gtg att aaa ggt ggc atg gtg gtc ttc tct 2064 Gly Val Lys Pro Lys Ile Val Ile Lys Gly Gly Met Val Val Phe Ser 670 675 680

gaa atg ggc gat tct aac gcg tcc gtg ccc acg cct cag ccg gtt tat 2112 Glu Met Gly Asp Ser Asn Ala Ser Val Pro Thr Pro Gln Pro Val Tyr 685 690 695 700

tac cgc gaa atg ttt ggg cac cac ggc aag gcg aaa ttt gac acc agc 2160

Tyr Arg Glu Met Phe Gly His His Gly Lys Ala Lys Phe Asp Thr Ser

705 710 715

atc act ttt cgt gtc tca agc gg 2183

Ile Thr Phe Arg Val Ser Ser
720

<210> 5

<211> 226

<212> PRT

<213> Helicobacter felis

<400> 5

Val Lys Leu Thr Pro Lys Glu Gln Glu Lys Phe Leu Leu Tyr Tyr Ala 1 5 10 15

Gly Glu Val Ala Arg Lys Arg Lys Ala Glu Gly Leu Lys Leu Asn Gln
20 25 30

Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala Arg 35 40 45

Arg Gly Lys Lys Thr Val Ala Glu Leu Met Glu Glu Cys Met His Phe 50 55 60

Leu Lys Lys Asp Glu Val Met Pro Gly Val Gly Asn Met Val Pro Asp 65 70 75 80

Leu Gly Val Glu Ala Thr Phe Pro Asp Gly Thr Lys Leu Val Thr Val 85 90 95

Asn Trp Pro Ile Glu Pro Asp Glu His Phe Lys Ala Gly Glu Val Lys

100	105	110

Phe Gly Cys Asp Lys Asp Ile Glu Leu Asn Val Gly Lys Glu Val Thr
115 120 125

Glu Leu Glu Val Thr Asn Glu Gly Pro Lys Ser Leu His Val Gly Ser 130 · 135 140

His Phe His Phe Phe Glu Thr Asn Lys Ala Leu Lys Phe Asp Arg Glu 145 150 155 160

Lys Ala Tyr Gly Lys Arg Leu Asp Ile Pro Ser Gly Asn Thr Leu Arg 165 170 175

Ile Gly Ala Gly Gln Thr Arg Lys Val Gln Leu Ile Pro Leu Gly Gly
180 185 190

Ser Lys Lys Val Ile Gly Met Asn Gly Leu Val Asn Asn Ile Ala Asp 195 200 205

Glu Arg His Lys His Lys Ala Leu Asp Lys Ala Lys Ser His Gly Phe 210 215 220

Ile Lys 225

<210> 6

<211> 496

<212> PRT

<213> Helicobacter felis

<400> 6

Met Lys Met Lys Lys Gln Glu Tyr Val Asn Thr Tyr Gly Pro Thr Thr

1 5 10 15

Gly Asp Lys Val Arg Leu Gly Asp Thr Asp Leu Trp Ala Glu Val Glu 20 25 30

His Asp Tyr Thr Thr Tyr Gly Glu Glu Leu Lys Phe Gly Ala Gly Lys
35 40 45

Thr Ile Arg Glu Gly Met Gly Gln Ser Asn Ser Pro Asp Glu Asn Thr 50 55 60

Leu Asp Leu Val Ile Thr Asn Ala Met Ile Ile Asp Tyr Thr Gly Ile
65 70 75 80

Tyr	Lys	Ala	Asp	Ile	Gly	Ile	Lys	Asn	GTÅ	гàг	шe	HIS	GTA	TTE	GIA
				85					90					95	

- Lys Ala Gly Asn Lys Asp Met Gln Asp Gly Val Ser Pro His Met Val 100 105 110
- Val Gly Val Gly Thr Glu Ala Leu Ala Gly Glu Gly Met Ile Ile Thr 115 120 125
- Ala Gly Gly Ile Asp Ser His Thr His Phe Leu Ser Pro Gln Gln Phe 130 135 140
- Pro Thr Ala Leu Ala Asn Gly Val Thr Thr Met Phe Gly Gly Gly Thr 145 150 155 160
- Gly Pro Val Asp Gly Thr Asm Ala Thr Thr Ile Thr Pro Gly Lys Trp
 165 170 175
- Asn Leu His Arg Met Leu Arg Ala Ala Glu Glu Tyr Ser Met Asn Val 180 185 190
- Gly Phe Leu Gly Lys Gly Asn Ser Ser Ser Lys Lys Gln Leu Val Glu 195 200 205
- Gln Val Glu Ala Gly Ala Ile Gly Phe Lys Leu His Glu Asp Trp Gly 210 215 220
- Thr Thr Pro Ser Ala Ile Asp His Cys Leu Ser Val Ala Asp Glu Tyr 225 230 235 240
- Asp Val Gln Val Cys Ile His Thr Asp Thr Val Asn Glu Ala Gly Tyr 245 250 255
- Val Asp Asp Thr Leu Asn Ala Met Asn Gly Arg Ala Ile His Ala Tyr 260 265 270
- His Ile Glu Gly Ala Gly Gly Gly His Ser Pro Asp Val Ile Thr Met 275 280 285
- Ala Gly Glu Leu Asn Ile Leu Pro Ser Ser Thr Thr Pro Thr Ile Pro 290 295 300
- Tyr Thr Ile Asn Thr Val Ala Glu His Leu Asp Met Leu Met Thr Cys 305 310 315 320
- His His Leu Asp Lys Arg Ile Arg Glu Asp Leu Gln Phe Ser Gln Ser 325 330 335

```
Arg Ile Arg Pro Gly Ser Ile Ala Ala Glu Asp Val Leu His Asp Ile
340 345 350
```

- Gly Val Ile Ala Met Thr Ser Ser Asp Ser Gln Ala Met Gly Arg Ala 355 360 365
- Gly Glu Val Ile Pro Arg Thr Trp Gln Thr Ala Asp Lys Asn Lys Lys 370 375 380
- Glu Phe Gly Lys Leu Pro Glu Asp Gly Ala Asp Asn Asp Asn Phe Arg 385 390 395 400
- Ile Lys Arg Tyr Ile Ser Lys Tyr Thr Ile Asn Pro Ala Leu Thr His
 405 410 415
- Gly Val Ser Glu Tyr Ile Gly Ser Val Glu Glu Gly Lys Ile Ala Asp 420 425 430
- Leu Val Val Trp Asn Pro Ala Phe Phe Gly Val Lys Pro Lys Ile Val 435 440 445
- Ile Lys Gly Gly Met Val Val Phe Ser Glu Met Gly Asp Ser Asn Ala 450 455 460
- Ser Val Pro Thr Pro Gln Pro Val Tyr Tyr Arg Glu Met Phe Gly His 465 470 475 480
- His Gly Lys Ala Lys Phe Asp Thr Ser Ile Thr Phe Arg Val Ser Ser 485 490 495
- <210> 7
- <211> 2883
- <212> DNA
- <213> Helicobacter felis
- <220>
- <221> CDS
- <222> (206)..(886)
- <220>
- <221> CDS
- <222> (897)..(2603)
- <400> 7
- rggragattt tecarcactt caageacata ttgateetgt gttgtgggtg gtaaattrer 60

Printed:10-07-2001


acti	tgtt	aat	rcta	ttat	ta a	tttt	ttaat	t aa	ttac	ttat	tate	cata	tat	aata	atatta	120
ttad	ctta	tat	taaa	aagt	ta a	taaa	aagta	a ac	gaaat	ttag	gact	tataa	atc	ccati	tgcctt	180
taaa	atti	taa	caca	agga	gt aa	atag								caa Gln	_	232
_		-	tta Leu					-		_	_	-	-		-	280
-	-		aag Lys					-	-		-			_	-	328
			gac Asp 45									_	_	_		376
_	_	-	tgc Cys				-			_	_	-	_			424
-			atg Met			-			_	_	_				-	472
	-		ctt Leu								-		-	-		520
		-	Gly						-	_		-				568
	-	-	aaa Lys 125											-		616
			cat His									-	-		_	ઈઈ4
-			ttc Phe	_	_	_		-				-		-		712







Ser	ggc Gly							Gln				760
	att Ile						Val				Gly	808
	aat Asn											856
	aaa Lys 220					taa	ggaq	gacto	_	Lys N	_	905
	caa Gln											953
	tta Leu											1001
	tat Tyr 265											1049
	atg Met											1097
	act Thr										_	1145
	G] À ààà	Ile										1193
	gac Asp											1241
Thr	gaa Glu . 345			G1y				Ile				1289







-		cac His									-	1337
		aca Thr 380	_						- •		-	1385
		act Thr								_		1433
		gca Ala								Phe	-	1481
		tct Ser					-	-		_	_	1529
		ttt Phe	_		_	-		-				1577
		tgc Cys 460										1625
	His	gat Asp									_	1673
		aac Asn	 _	_			_					1721
		cac His										1769
		tcc Ser										1817
		cac His 540		_		-		_				1865

		atc Ile	Glu			Ser		_		Arg	1913
		atc Ile 570									1961
		agc Ser							-		2009
		act Thr						-			2057
		gaa Glu							-	-	2105
		aaa Lys								_	2153
		ggc Gly 650					-	_	-		2201
		gcc Ala									2249
		gtc Val									2297
		ccg Pro								-	2345
		gac Asp								-	2393
	Val	aaa Lys 730							_	-	2441



002025(55.3(1/10)/2000)

Lys		•	-				_		-		-			gac Asp		2489
-					_	_	_					_		gta Val	-	25 <u>,</u> 37
			•						_					gcc Ala	-	2585
-		act Thr			tag	gcad	aato	jcc (cctt	tgg	là đợ	aggt	tat	t		2633
ttag	gaat	ct t	cato	caaac	eg ca	cctç	gcaat	c cg	gtctt	gcg	tgto	gcgat	cg	tgtcg	gcttta	2693
aaac	aact	tt t	cato	ttta	aa go	aato	gcca	a ttt	ttaa	itta	ättt	aatt	ct	tataa	attaat	2753
atta	tati	at q	jeced	ectca	at t	ttaa	agga	a gaa	ttat	gcg	tagç	gtctt	tg (gtatt	gctat	2813
gtgg	ıgggt	tg t	ttg	gtgct	g gg	jcgca	aagg	g gta	ttga	aac	ccat	cgcc	etc a	aaaaa	agtag	2873
aago	caca	agg										,				2883
<210 <211 <212 <213	.> 22 !> PI		obact	er 1	felis	5										
<211 <212	.> 22 !> PI !> He	۲r	bbact	ter 1	felis	5										
<211 <212 <213 <400	> 22 2> PI 3> He 0> 8	RT elico					Gln	Glu	Lys 10	Phe	Leu	Leu	Tyr	Туг 15	Ala	
<211 <212 <213 <400 Val	.> 22 !> PH !> He !> 8 Lys	RT elico Leu	Thr	Pro 5	Lys	Glu			10					15 Asn		
<211 <212 <213 <400 Val 1	> 22 > PR 3> He 0> 8 Lys	RT elico Leu Val	Thr Ala 20	Pro 5 Arg	Lys Lys	Glu Arg	Lys	Ala 25	10 Glu	Gly	Leu	Lys	Leu 30	15 Asn	Gln	
<211 <212 <213 <400 Val 1 Gly Pro	.> 22 ?> PR ?> He ?> 8 Lys Glu	Leu Val Ala 35	Thr Ala 20 Ile	Pro 5 Arg	Lys Lys Tyr	Glu Arg Ile	Lys Ser 40	Ala 25 Ala	10 Glu His	Gly Ile	Leu Met	Lys Asp 45	Leu 30 Glu	15 Asn	Gln Arg	





Leu Gly Val Glu Ala Thr Phe Pro Asp Gly Thr Lys Leu Val Thr Val 85 90 Asn Trp Pro Ile Glu Pro Asp Glu His Phe Lys Ala Gly Glu Val Lys 100 105 Phe Gly Cys Asp Lys Asp Ile Glu Leu Asn Ala Gly Lys Glu Val Thr 115 120 Glu Leu Glu Val Thr Asn Glu Gly Pro Lys Ser Leu His Val Gly Ser 135 His Phe His Phe Phe Glu Ala Asn Lys Ala Leu Lys Phe Asp Arg Glu 150 155 Lys Ala Tyr Gly Lys Arg Leu Asp Ile Pro Ser Gly Asn Thr Leu Arg 165 170 Ile Gly Ala Gly Gln Thr Arg Lys Val Gln Leu Ile Pro Leu Gly Gly 180 185 190 Ser Lys Lys Val Ile Gly Met Asn Gly Leu Val Asn Asn Ile Ala Asp 195 200 205 Glu Arg His Lys His Lys Ala Leu Asp Lys Ala Lys Ser His Gly Phe 210 215 220 Ile Lys 225 <210> 9 <211> 568 <212> PRT <213> Helicobacter felis Met Lys Met Lys Gln Glu Tyr Val Asn Thr Tyr Gly Pro Thr Lys 1 5 15 Gly Asp Lys Val Arg Leu Gly Asp Thr Asp Leu Trp Ala Glu Val Glu 20 25 His Asp Tyr Thr Thr Tyr Gly Glu Leu Lys Phe Gly Ala Gly Lys 40 .

Thr Ile Arg Glu Gly Met Gly Gln Ser Asn Ser Pro Asp Glu Asn Thr 50 55 60

-	гте
65 70 75	80

- Tyr Lys Ala Asp Ile Gly Ile Lys Asn Gly Lys Ile His Gly Ile Gly
 85 90 95
- Lys Ala Gly Asn Lys Asp Met Gln Asp Gly Val Ser Pro His Met Val 100 105 110
- Val Gly Val Gly Thr Glu Ala Leu Ala Gly Glu Gly Met Ile Ile Thr 115 120 125
- Ala Gly Gly Ile Asp Ser His Thr His Phe Leu Ser Pro Gln Gln Phe 130 135 140
- Pro Thr Ala Leu Ala Asn Gly Val Thr Thr Met Phe Gly Gly Gly Thr 145 150 155 160
- Gly Pro Val Asp Gly Thr Asn Ala Thr Thr Ile Thr Pro Gly Lys Trp 165 170 175
- Asn Leu His Arg Met Leu Arg Ala Ala Glu Glu Tyr Ser Met Asn Val 180 185 190
- Gly Phe Leu Gly Lys Gly Asn Ser Ser Ser Lys Lys Gln Leu Val Glu 195 200 205
- Gln Val Glu Ala Gly Ala Ile Gly Phe Lys Leu His Glu Asp Trp Gly 210 215 220
- Thr Thr Pro Ser Ala Ile Asp His Cys Leu Ser Val Ala Asp Glu Tyr 225 230 235 240
- Asp Val Gln Val Cys Ile His Thr Asp Thr Val Asn Glu Ala Gly Tyr 245 250 255
- Val Asp Asp Thr Leu Asn Ala Met Asn Gly Arg Ala Ile His Ala Tyr 260 265 270
- His Ile Glu Gly Ala Gly Gly His Ser Pro Asp Val Ile Thr Met 275 280 285
- Ala Gly Glu Leu Asn Ile Leu Pro Ser Ser Thr Thr Pro Thr Ile Pro 290 295 300
- Tyr Thr Ile Asn Thr Val Ala Glu His Leu Asp Met Leu Met Thr Cys 305 310 315 320



His	His	Leu	Asp	Lys	Arg	Ile	Arg	Glu	Asp	Leu	Gln	Phe	Ser	Gln	ser
				325					330					335	


- Arg Ile Arg Pro Gly Ser Ile Ala Ala Glu Asp Val Leu His Asp Met 340 345 350
- Gly Val Ile Ala Met Thr Ser Ser Asp Ser Gln Ala Met Gly Arg Ala 355 360 365
- Gly Glu Val Ile Pro Arg Thr Trp Gln Thr Ala Asp Lys Asn Lys Lys 370 375 380
- Glu Phe Gly Lys Leu Pro Glu Asp Gly Lys Asp Asn Asp Asn Phe Arg 385 390 395 400
- Ile Lys Arg Tyr Ile Ser Lys Tyr Thr Ile Asn Pro Ala Leu Thr His
 405 410 415
- Gly Val Ser Glu Tyr Ile Gly Ser Val Glu Glu Gly Lys Ile Ala Asp 420 425 430
- Leu Val Val Trp Asn Pro Ala Phe Phe Gly Val Lys Pro Lys Ile Val 435 440 445
- Ile Lys Gly Gly Met Val Val Phe Ser Glu Met Gly Asp Ser Asn Ala 450 455 460
- Ser Val Pro Thr Pro Gln Pro Val Tyr Tyr Arg Glu Met Phe Gly His 465 470 475 480
- His Gly Lys Ala Lys Phe Asp Thr Ser Ile Thr Phe Val Ser Lys Val 485 490 495
- Ala Tyr Glu Asn Gly Val Lys Glu Lys Leu Gly Leu Glu Arg Gln Val 500 505 510
- Leu Pro Val Lys Asn Cys Arg Asn Ile Thr Lys Lys Asp Phe Lys Phe 515 520 525
- Asn Asp Lys Thr Ala Lys Ile Thr Val Asp Pro Lys Thr Phe Glu Val 530 535 540
- Phe Val Asp Gly Lys Leu Cys Thr Ser Lys Pro Thr Ser Gln Val Pro 545 550 560

Leu Ala Gln Arg Tyr Thr Phe Phe 565

<21	0> 1	0														
<21	1> 2	407								•						
<21	2> DI	AN														
<21	3> H	elic	obac	ter	feli	s							•			
<22	0>															
<22	l> CI	os														
<222	2> (2	2)	(682)												
<220																
	L> CI															
<222	2> ((593)	(2	399)												
-10	N 1/	`														
	0> 10		ta 2	a							+1					40
						_	_	-		-		_			at gcg yr Ala	49
V 6	1	An m	eu II	IIL P.	Б	ys G.	iu G.	111 G.	•	ys P 10	.16 T	su b	eu I	•	yr Ala 15	
	•				,					10					10	
aac	gaa	ata	act	aσa	aaσ	cac	aaa	aca	σασ	aac	tta	aaσ	ctc	aac	caa	97
	-		-	_	_	-						-		Asn		
_			20	-	•	,	•	25		•		•	30			
ccc	gaa	gcc	att	gcc	tac	att	agt	gcc	cat	att	atg	gac	gag	gcg	cgc	145
Pro	Glu	Ala	Ile	Ala	Tyr	Ile	Ser	Ala	His	Ile	Met	Asp	Glu	Ala	Arg	
		35					40					45				
cgt	ggc	aaa	aag	acc	gtt	gcg	gaa	ctt	atg	gaa	gag	tgt	atg	cac	ttt	193
Arg	Gly	ŗys	Lys	Thr	Val	Ala	Glu	Leu	Met	Glu	Glu	Cys	Met	His	Phe	
	50					55					60			-		
			-			_						-		cct		24:
	Lys	Lys	Asp	GLu		Met	Pro	GLA	Val	_	Asn	Met	Val	Pro		
65					70					75					80	
tta	aac	ata	gaa	act	act	++ +	CCC.	rat	aac	acc	aaa	ctc	αta	acc	ata	289
								_						Thr		
				85					90		-,-			95		
aat	tgg	ccc	atc	gaa	ccc	gat	qaa	cac	ttc	aaa	gcg	ggc	gaa	gtc	aaa	33
Asn	Trp	Pro	Ile	Glu	Pro	Asp	Glu	His	Phe	Lys	Ala	Gly	Glu	Val	Lys	
	-		100			=		105					110			
ttt	ggc	tgt	gat	aaa	gac	att	gaa	ctc	aac	gca	ggt	aag	gaa	gtt	acc	385
Phe	Gly	Cys	Asp	Lys	Asp	Ile	Glu	Leu	Asn	Ala	Gly	Lys	Glu	Val	Thr	
		115					120					125				

						gaa										433
Glu			Val	Thr	Asn	Glu		Pro	Lys	Ser			Val	Gly	Ser	
	130					135					140					
cat	ttc	cac	ttc	ttt	gaa	gcc	aac	aag	σса	tta	aaa	ttc	gat	caa	722	481
						Ala										401
145					150			_,_		155	-1-		p	7129	160	
						cta							_		_	529
Lys	Ala	Tyr	Gly			Leu	Asp	Ile			Gly	Asn	Thr	Leu	Arg	
				165					170					175		
att	aaa	qca	qqa	caa	acc	cgt	aaa	ata	cag	tta	atc	cct	ctt	aac	aac	577
						Arg										5,,
			180					185					190	-	•	
												•				
						atg								-	-	625
ser	гÀг	Lуs 195	vaı	TIE	GTA	Met	200	GIA	Leu	Val	Asn		Ile	Ala	Asp	
		100					200					205				
gaa	cgc	cat	aaa	cac	aaa	gcg	tta	gaa	aaa	gca	aaa	tct	cac	gga	ttt	673
Glu	Arg	His	Lys	His	Lys	Ala	Leu	Glu	Lys	Ala	Lys	Ser	His	Gly	Phe	
	210					215					220			-		
ata	222	taa	aa - a	acto		ıtg a		+~ ~				+				700
Ile		caa	ggag	accc		icy a let I							-			722
225	-		•					30		-7			35	u 1 1		
						ggc							-		-	770
Thr	Tyr	G1y 240	Pro	Thr	Thr	Gly		Lys	Val	Arg	Leu		Asp	Thr	Asp	
		240					245					250				
ctt	tgg	gca	gaa	gta	gaa	cat	gac	tat	acc	act	tat	ggc	gaa	gag	ctc	818
						His							-	-		
	255					260					265					
_																
						act										866
195 270	FIIC	сту	Ата		275	Thr	ire	Arg	GIU	280	met	етА	GIN		Aşn 285	
. , ,										200					203	
agt	cca	gat	gaa	aac	acc	cṫa	gat	tta	gtc	atc	acc	aac	gcg	atg	att	914
Ser	Pro	Asp	Glu .	Asn	Thr	Leu	Asp	Leu	Val	Ile	Thr .	Asn	Ala	Met	Ile	
				290					295					300		
·++	as.	tac	200	aaa	2++	tac	222	acc	~ ~ ~	a++	~~~	_+-				0.00
			qCC	999	act	cac	aad	gee	yac	alt	yyc	all	aaa	aat	qqc	962
TE	azA	Tvr	Thr	Glv	Ile	Tvr	Lvs	Ala	Asn		Glv	Tle	Lve	Den		
TIE	Asp		Thr 305	Gly	Ile	Tyr		Ala 310	Asp		Gly	Ile	Lys 315	Asn		

atc Ile									1010
agc Ser 335									1058
ggt Gly									1106
tct Ser									1154
ttt Phe								-	1202
act Thr							_	-	1250
tat Tyr 415									1298
aaa Lys									1346
cat His							_	-	1394
gta Val							_	_	1442
aat Asn									1490
gcc Ala 495									1538





	Asp				Ala				Ile			tcc Ser 525	1586
		act Thr		Pro				Thr				Leu	1634
		atg Met 545											1682
		tcc Ser											1730
		cat His											1778
		G] y											1826
		aat Asn											1874
		aac Asn 625								Lys			1922
		cta Leu			Gly							_	1970
Glu		atc Ile											2018
		aag Lys	Ile				Gly						2066
		tcc Ser											2114

cgc gaa atg ttt ggg cat cac ggc aag gcg aaa ttt gac acc agc at Arg Glu Met Phe Gly His His Gly Lys Ala Lys Phe Asp Thr Ser Il 705 710 715	
act ttt gtt tcc aaa gtc gcc tat gaa aat ggc gtg aaa gaa aaa ct Thr Phe Val Ser Lys Val Ala Tyr Glu Asn Gly Val Lys Glu Lys Le 720 725 730	
ggc tta gag cgc aag gtg cta ccc gtg aaa aac tgc cgc aac atc ac Gly Leu Glu Arg Lys Val Leu Pro Val Lys Asn Cys Arg Asn Ile Th 735 740 745	
aag aaa gac ttc aaa ttc aac aac aag acg gcg cat atc act gtc ga Lys Lys Asp Phe Lys Phe Asn Asn Lys Thr Ala His Ile Thr Val As 750 760 760	•
cct aaa acc ttc gag gtc ttt gta gat ggc aaa ctc tgc acc tct aac Pro Lys Thr Phe Glu Val Phe Val Asp Gly Lys Leu Cys Thr Ser Lys 770 775 780	
ccc gcc tct gaa gtg cct cta gcc cag cgc tac act ttc ttc tag Pro Ala Ser Glu Val Pro Leu Ala Gln Arg Tyr Thr Phe Phe 785 790 795	2399
gcncaatg	2407
<210> 11 <211> 226 <212> PRT <213> Helicobacter felis	
: <400> 11	
Val Lys Leu Thr Pro Lys Glu Gln Glu Lys Phe Leu Leu Tyr Tyr Al 1 5 10 15	1
Gly Glu Val Ala Arg Lys Arg Lys Ala Glu Gly Leu Lys Leu Asn Glu 20 25 30	ı
25 25	
Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala Arc 35 40 45	3
Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala Arc	



Leu Gly Val Glu Ala Thr Phe Pro Asp Gly Thr Lys Leu Val Thr Val 85 90 95

Asn Trp Pro Ile Glu Pro Asp Glu His Phe Lys Ala Gly Glu Val Lys
100 105 110

Phe Gly Cys Asp Lys Asp Ile Glu Leu Asn Ala Gly Lys Glu Val Thr 115 120 125

Glu Leu Glu Val Thr Asn Glu Gly Pro Lys Ser Leu His Val Gly Ser 130 135 140

Lys Ala Tyr Gly Lys Arg Leu Asp Ile Pro Ser Gly Asn Thr Leu Arg 165 170 175

Ile Gly Ala Gly Gln Thr Arg Lys Val Gln Leu Ile Pro Leu Gly Gly
180 185 190

Ser Lys Val Ile Gly Met Asn Gly Leu Val Asn Asn Ile Ala Asp 195 200 205

Glu Arg His Lys His Lys Ala Leu Glu Lys Ala Lys Ser His Gly Phe 210 215 220

Ile Lys

225

<210> 12

<211> 568

<212> PRT

<213> Helicobacter felis

<400> 12

Met Lys Met Lys Lys Gln Glu Tyr Val Asn Thr Tyr Gly Pro Thr Thr 1 5 10 15

Gly Asp Lys Val Arg Leu Gly Asp Thr Asp Leu Trp Ala Glu Val Glu
20 25 30

His Asp Tyr Thr Thr Tyr Gly Glu Glu Leu Lys Phe Gly Ala Gly Lys
35 40 45


Thr Ile Arg Glu Gly Met Gly Gln Ser Asn Ser Pro Asp Glu Asn Thr

•	^
	"

- Leu Asp Leu Val Ile Thr Asn Ala Met Ile Ile Asp Tyr Thr Gly Ile
 65 75 80
- Tyr Lys Ala Asp Ile Gly Ile Lys Asn Gly Lys Ile His Gly Ile Gly 85 90 95
- Lys Ala Gly Asn Lys Asp Met Gln Asp Gly Val Ser Pro His Met Val
 100 105 110
- Val Gly Val Gly Thr Glu Ala Leu Ala Gly Glu Gly Met Ile Ile Thr 115 120 125
- Ala Gly Gly Ile Asp Ser His Thr His Phe Leu Ser Pro Gln Gln Phe 130 135 140
- Pro Thr Ala Leu Ala Asn Gly Val Thr Thr Met Phe Gly Gly Gly Thr 145 150 155 160
- Gly Pro Val Asp Gly Thr Asn Ala Thr Thr Ile Thr Pro Gly Lys Trp
 165 170 175
- Asn Leu His Arg Met Leu Arg Ala Ala Glu Glu Tyr Ser Met Asn Val 180 185 190
- Gly Phe Leu Gly Lys Gly Asn Ser Ser Ser Lys Lys Gln Leu Val Glu 195 200 205
- Gln Ile Glu Ala Gly Ala Ile Gly Phe Lys Leu His Glu Asp Trp Gly 210 215 220
- Thr Thr Pro Ser Ala Ile Asp His Cys Leu Ser Val Ala Asp Glu Tyr 225 230 235 240
- Asp Val Gln Val Cys Ile His Thr Asp Thr Val Asn Glu Ala Gly Tyr 245 250 255
- Val Asp Asp Thr Leu Asn Ala Met Asn Gly Arg Ala Ile His Ala Tyr 260 265 270
- His Ile Glu Gly Ala Gly Gly Gly His Ser Pro Asp Val Ile Thr Met 275 280 . 285
- Ala Gly Glu Leu Asn Ile Leu Pro Ser Ser Thr Thr Pro Thr Ile Pro 290 295 300
- Tyr Thr Ile Asn Thr Val Ala Glu His Leu Asp Met Leu Met Thr Cys

305	i				310					315					320
His	His	Leu	Asp	Lys 325		Ile	Arg	Glu	Asp 330		Gln	Phe	Ser	Gln 335	
Arg	Ile	Arg	Pro 340	Gly	Ser	Ile	Ala	Ala 345		Asp	Val	Leu	His 350	-	Ile
Gly	Val	Ile 355		Met	Thr	Ser	Ser 360	Asp	Ser	Gln	Ala	Met 365	Gly	Arg	Ala
Gly	Glu 370	Val	Ile	Pro	Arg	Thr 375	Trp	Gln	Thr	Ala	Asp 380		Asn	Lys	Lys
Glu 385	Phe	Gly	Lys	Leu	Pro 390	Glu	Asp	Ser	Ala	Asp 395	Asn	Asp	Asn	Phe	Arg 400
Ile	Lys	Arg	Tyr	Ile 405	Ser	Lys	Tyr	Thr	Ile 410		Pro	Ala	Leu	Thr 415	His
Gly	Val	Ser	Glu 420	Tyr	Ile	Gly	Ser	Val 425	Glu	Glu	Gly	Lys	Ile 430	Ala	Asp
Leu	Val	Val 435	Trp	Asn	Pro	Ala	Phe 440	Phe	Gly	Val	Lys	Pro 445	Lys	Ile	Val
Ile	Lys 450	Gly	Gly	Met	Val	Val 455	Phe	Ser	Glu	Met	Gly 460	Asp	Ser	Asn	Ala
Ser 465	Val	Pro	Thr	Pro	Gln 470	Pro	Val	Tyr	Tyr	Arg 475	Glu	Met	Phe	Gly	His 480
His	Gly	Lys	Ala	Lys 485	Phe	Asp	Thr	Ser	Ile 490	Thr	Phe	Val	Ser	Lys 495	Val
Ala	Tyr	Glu	Asn 500	Gly	Val	Lys	Glu	Lys 505	Leu	Gly	Leu	Glu	Arg 510	Lys	Val
		515					520					525	Phe		
	530					535					540		Phe-		
Phe 545	Val	Asp	Gly	Lys	Leu 550	Cys	Thr	Ser	Lys	Pro 555	Ala	Ser	Glu	Val	Pro 560

Leu Ala Gln Arg Tyr Thr Phe Phe



<21	0> 1	3								•						
<21	1> 2	452														
<21	2> D	NA														
<21	3> H	elic	obac	ter	feli	s										
<22	0>															
<22	1> C	DS														
<222	2> (48).	. (72	8)												
<22																
	1> C															
<222	2> (739)	(2	445)											•	
-401	0> 1:	2														
_	_		tece	atta	cc ti	ttaaa	aatti	t aad	racaa	anna	otaa	atad	ata	222	ctc	56
499				9					Juou	-99-	g _i cu,	acuy		Lys		30
													1	-,-		
aca	ccc	aaa	gag	caa	gaa	aag	ttc	ttg	tta	tat	tat	gcg	ggc	gaa	gtg	104
Thr	Pro	Lys	Glu	Gln	Glu	Lys	Phe	Leu	Leu	Tyr	Tyr	Ala	Gly	Glu	Val	
	5					10					15					
gct	aga	aag	cgc	aaa	gca	gag	ggc	tta	aag	ctc	aac	caa	ccc	gaa	gcc	152
	Arg	Lys	Arg	Lys	Ala	Glu	Gly	Leu	Lys	Leu	Asn	Gln	Pro	Glu	Ala	
20					25				•	30					35	
						+										200
	_			-	gcc			_	-			-	-			200
TIG	ALA	TAT	TIE	40	Ala	UIS	тте	Met	ASP 45	GIU	АТА	Arg	Arg	50	_	
•		-		40					43							
aaa	acc	qtt	aca	qaa	ctt	atq	gaa	qaq	tqt	atq	cac	ttt	ttq	aaa	aaa	248
		-		_	Leu	_	-		-	_			_			
-			55					60					65	_	_	
gac	gag	gtg	atg	ccc	ggg	gtg	ggg	aat	atg	gtc	cct	gat	ttg	ggc	gţg	296
Asp	Glu	Val	Met	Pro	Gly	Val	Gly	Asn	Met	Val	Pro	Asp	Leu	Gly	Val	
		70					75					80				
						•										
-	-				gat -					-						344
GIu		Thr	Pne	Pro	Asp	_	Thr	. Lys	Leu	Val		val	Asn	Trp	Pro	
	85					90					95					
atc	gaa	cct	gat	gaa	cac	ttt	aaσ	aca	aat	gaa	ata	aaa	ttt	aac	tat	392
	_		-		His		-			-					-	

100)		•		105			110					115	
		gac Asp			Leu			Glu					Glu	440
		aac Asn												488
		gaa Glu 150								_		-		536
		cgc Arg											-	584
		acc Thr									_			632
		ggc Gly												680
		aaa Lys										_	taa	728
ggaç	gacto	ecc a		ys M			lu T					yr (-	777
		aca Thr											-	825
		gaa Glu												873
		aaa Lys 275												921
		acc Thr						-	_			-		969

290 2	95 3	00
-------	------	----

													•			
acc	ggg	att	tac	aaa	gcc	gac	att	ggc	att	aaa	aat	ggc	aaa	atc	cat	1017
Thr	Gly	Ile	Tyr	Lys	Ala	Asp	Ile	Gly	Ile	Lys	Asn	Gly	Lys	Ile	His	
305	_		-	•	310	•		-		315		-	•	·	320	
															520	
aac	a++	aac	224	a a.	<i>aa</i> ₂	226	224	~ 2 ~	2+4	C 222	~-+	~~~	~+~	200		1065
			_		gga		_	-	-		-		-	-		1065
GIY	тте	GIŸ	rys		Gly	Asn	гÀг	Asp			Asp	GIÀ	Val		Pro	
				325					330					335		
cat	atg	gtc	gtg	ggt	gtg	ggc	aca	gaa	gca	cta	gca	ggg	gaa	ggt	atg	1113
His	Met	Val	Val	Gly	Val	Gly	Thr	Glu	Ala	Leu	Ala	Gly	Glu	Gly	Met	
		•	340					345					350			
att	att	acc	gct	ggg	ggg	atc	gat	tca	cac	acc	cac	ttc	ctc	tct	cca	1161
					Gly		-									
		355		-	•		360					365				
C22	caa	ttc	cct	200	gct	cta	~~~	22+	~~~	~++	202	202	2 t &	+++	~~~	1209
										_			_			1209
GIII		Pile	PIO	THE	Ala		AId	ASII	GTA	Val		THE	мес	Pne	стА	
	370					375					380					
			_		gta	_		_							_	1257
Gly	Gly	Thr	Gly	Pro	Val	Asp	Gly	Thr	Asn	Ala	Thr	Thr	Ile	Thr	Pro	
385					390					395					400	
ggc	aaa	tgg	aac	ttg	cac	cgc	atg	ttg	cgc	gca	gca	gaa	gag	tat	tct	1305
Gly	Lys	Trp	Asn	Leu	His	Arg	Met	Leu	Arg	Ala	Ala	Glu	Glu	Tyr	Ser	
	•			405					410					415		
ato	aat	ata	aac	ttt	ttg	aac	aaa	aac	aat	agc	tct	agt	aaa	aaa	caa	1353
_					Leu					-		_				
			420			,	~, ~	425	- 1011				430	_,_	01	
			120					120					100			
a++	ata	~ ~~	G 2 2	at a	~~~	~~~	~~~	~~~	-++	~~+			++~		~~~	1401
	-	-		_	gaa								_		-	1401
Leu	vaı		GIN	vaı	Glu	Ата	_	Ala	TTE	сту	Pne	-	Leu	HIS	GIU	
		435					440					445				
gac	tgg	ggc	aca	act	cca	agt	gcg	atc	gat	cac	tgc	ttg	agc	gta	gca	1449
Asp	Trp	Gly	Thr	Thr	Pro	Ser	Ala	Ile	Asp	His	Cys	Leu	Ser	Val	Ala	
	450					455					460					
gat	gaa	tac	gat	gtg	caa	gtt	tgt	ata	cac	acc	gat	acg	gtc	aat	gag	1497
Asp	Glu	Tyr	Asp	Val	Gln	Val	Cys	Ile	His	Thr	Asp	Thr	Val	Asn	Glu	
465		-	-		470		-			475	-				480	
gca	aat	tat	gta	gat	gac	acc	cta	aat	σca	atσ	aac	ggg	cac	acc	atc	1545
_					Asp				_	_			-	-		
	y	- 2.		12			u	11	4			×	4			

485 490 495

cat	gcc	tac	cac	att	gag	gga	gcg	ggt	gga	gga	cac	tca	cct	gat	qtt	1593
His	Ala	Tyr	His	Ile	Glu	Gly	Ala	Gly	Gly	Gly	His	Ser	Pro	Asp	Val	
			500					505		•			510	_		
atc	acc	atg	gca	ggc	gaa	gtg	aat	att	cta	ccc	tcc	tcc	aca	acc	cct	1641
Ile	Thr	Met	Ala	Gly	Glu	Val	Asn	Ile	Leu	Pro	Ser	Ser	Thr	Thr	Pro	
		515					520					525				
act	atc	ccc	tat	acc	att	aat	acg	gtt	gca	gaa	cac	tta	gac	atg	ctt	1689
Thr	Ile	Pro	Tyr	Thr	Ile	Asn	Thr	Val	Ala	Glu	His	Leu	Asp	Met	Leu	
	530					535					540		_			
atg	acc	tgc	cac	cac	cta	gat	aaa	cgc	atc	cgc	gag	gat	ctc	caa	ttt	1737
Met	Thr	Cys	His	His	Leu	Asp	Lys	Arg	Ile	Arg	Glu	Asp	Leu	Gln	Phe	
545					550					555					560	
							•									
tct	caa	agc	cgt	atc	cgc	ccc	ggc	tct	atc	gcc	gct	gaa	gat	gtg	ctc	1785
Ser	Gln	Ser	Arg	Ile	Arg	Pro	Gly	Ser	Ile	Ala	Ala	Glu	Asp	Val	Leu	
				565					570					575		
cat	gat	atc	ggt	gtg	atc	gcg	atg	aca	agt	tcc	gat	tcg	caa	gca	atg	1833
His	Asp	Ile	Gly	Val	Ile	Ala	Met	Thr	Ser	Ser	Asp	Ser	Gln	Ala	Met	
			580					585					590			
		gct											-	-	-	1881
Gly	Arg	Ala	Gly	Glu	Val	Ile	Pro	Arg	Thr	Trp	Gln	Thr	Ala	Asp	Lys	
		595					600					605				
		aaa									-		_		_	1929
Asn	Lys	Lys	Glu	Phe	Gly	Lys	Leu	Pro	Glu	Asp	Gly	Ala	Asp	Asn	Asp	
	610					615					620					
												•				
		cgc													-	1977
	Phe	Arg	Ile	Lys		Tyr	Ile	Ser	Lys		Thr	Ile	Asn	Pro	Ala	
625					630					635					640	
		cat														2025
Leu	Thr	His	Gly		Ser	Glu	Tyr	Ile	-	Ser	Val	Glu	Glu	-	Lys	
				645					650					655		
		gac -														2073 [.]
Ile	Ala	Asp		val	val	Trp	Asn		Ala	Phe	Phe	Gly		Lys	Pro	
			660					665					670			
		gtg														2121
Lys	Ile	Val	Ile	Lys	Gly	Gly	Met	Val	Val	Phe	Ser	Glu	Met	Gly	Asp	*





. _ .										~++		.		~~~		2160
				gtg												2169
ser		ATA	ser	Val	Pro		Pro	GIN	Pro	vai	_	ryr	Arg	GIU	met	
	690					695					700					
				ggc	_	,			_		-				•	2217
	Gly	His	His	Gly	_	Ala	Lys	Phe	Asp		Ser	Ile	Thr	Phe		
705					710					715					720	
		-	_	tat	_					_						2265
Ser	Lys	Val	Ala	Tyr	Glu	Asn	Gly	Val	_	Glu	Lys	Leu	Gly			
				725					730					735		
_	-			ccc				-	-				_	_		2313
Arg	Lys	Val		Pro	Val	Lys	Asn	-	Arg	Asn	Ile	Thr	-	Lys	Asp	
			740					745					750			
	_			gac			-				-	_	-			2361
Phe	Lys	Phe	Asn	Asp	Lys	Thr	Ala	Lys	Ile	Thr	Val	_	Pro	Lys	Thr	
		755					760					765				
		_		gta	_				_							2409
Phe		Val	Phe	Val	Asp	_	Lys	Leu	Cys	Thr		Lys	Pro	Thr	Ser	
	770					775					780					
-	-			gcc							tag	gcat	taat			2452
	Val	Pro	Leu	Ala		Arg	Tyr	Thr	Phe							
785					790					795						
											٠					
_	<210> 14															
	<211> 226															
_	2> PI		a bi a		6 01÷	_										
<21:	52 HE	STIC	Dac	ter :	гетт	5										

<213> Helicobacter felis

<400> 14

Val Lys Leu Thr Pro Lys Glu Gln Glu Lys Phe Leu Leu Tyr Tyr Ala

Gly Glu Val Ala Arg Lys Arg Lys Ala Glu Gly Leu Lys Leu Asn Gln

Pro Glu Ala Ile Ala Tyr Ile Ser Ala His Ile Met Asp Glu Ala Arg

Arg Gly Lys Lys Thr Val Ala Glu Leu Met Glu Glu Cys Met His Phe

Leu Lys Lys Asp Glu Val Met Pro Gly Val Gly Asn Met Val Pro Asp 65 70 75 80

Leu Gly Val Glu Ala Thr Phe Pro Asp Gly Thr Lys Leu Val Thr Val 85 90 95

Asn Trp Pro Ile Glu Pro Asp Glu His Phe Lys Ala Gly Glu Val Lys 100 105 110

Phe Gly Cys Asp Lys Asp Ile Glu Leu Asn Ala Gly Lys Glu Val Thr 115 120 125

Glu Leu Glu Val Thr Asn Glu Gly Pro Lys Ser Leu His Val Gly Ser 130 135 140

His Phe His Phe Phe Glu Ala Asn Lys Ala Leu Lys Phe Asp Arg Glu 145 150 155 160

Lys Ala Tyr Gly Lys Arg Leu Asp Ile Pro Ser Gly Asn Thr Leu Arg 165 170 175

Ile Gly Ala Gly Gln Thr Arg Lys Val Gln Leu Ile Pro Leu Gly Gly 180 185 190

Ser Lys Lys Val Ile Gly Met Asn Gly Leu Val Asn Asn Ile Ala Asp 195 200 205

Glu Arg His Lys His Lys Ala Leu Asp Lys Ala Lys Ser His Gly Phe 210 215 220

Ile Lys

225

<210> 15

<211> 568

<212> PRT

<213> Helicobacter felis

<400> 15

Met Lys Met Lys Lys Gln Glu Tyr Val Asn Thr Tyr Gly Pro Thr Thr 1 5 10 15

Gly Asp Lys Val Arg Leu Gly Asp Thr Asp Leu Trp Ala Glu Val Glu 20 25 30

His Asp Tyr Thr Thr Tyr Gly Glu Glu Leu Lys Phe Gly Ala Gly Lys

3			
•	~		

- Thr Ile Arg Glu Gly Met Gly Gln Ser Asn Ser Pro Asp Glu Asn Thr
 50 55 60
- Leu Asp Leu Val Ile Thr Asn Ala Met Ile Ile Asp Tyr Thr Gly Ile
 65 70 75 80
- Tyr Lys Ala Asp Ile Gly Ile Lys Asn Gly Lys Ile His Gly Ile Gly
 85 90 95
- Lys Ala Gly Asn Lys Asp Met Gln Asp Gly Val Ser Pro His Met Val
 100 105 110
- Val Gly Val Gly Thr Glu Ala Leu Ala Gly Glu Gly Met Ile Ile Thr 115 120 125
- Ala Gly Gly Ile Asp Ser His Thr His Phe Leu Ser Pro Gln Gln Phe 130 135 140
- Pro Thr Ala Leu Ala Asn Gly Val Thr Thr Met Phe Gly Gly Gly Thr 145 150 155 160
- Gly Pro Val Asp Gly Thr Asn Ala Thr Thr Ile Thr Pro Gly Lys Trp
 165 170 175
- Asn Leu His Arg Met Leu Arg Ala Ala Glu Glu Tyr Ser Met Asn Val 180 185 190
- Gly Phe Leu Gly Lys Gly Asn Ser Ser Ser Lys Lys Gln Leu Val Glu 195 200 205
- Gln Val Glu Ala Gly Ala Ile Gly Phe Lys Leu His Glu Asp Trp Gly 210 215 220
- Thr Thr Pro Ser Ala Ile Asp His Cys Leu Ser Val Ala Asp Glu Tyr 225 230 235 240
- Asp Val Gln Val Cys Ile His Thr Asp Thr Val Asn Glu Ala Gly Tyr 245 250 255
- Val Asp Asp Thr Leu Asn Ala Met Asn Gly Arg Ala Ile His Ala Tyr 260 265 270
- His Ile Glu Gly Ala Gly Gly Gly His Ser Pro Asp Val Ile Thr Met 275 280 285
- Ala Gly Glu Val Asn Ile Leu Pro Ser Ser Thr Thr Pro Thr Ile Pro

290	205	2	^ ^
230	295	3	00

- Tyr Thr Ile Asn Thr Val Ala Glu His Leu Asp Met Leu Met Thr Cys 305 310 315 320
- His His Leu Asp Lys Arg Ile Arg Glu Asp Leu Gln Phe Ser Gln Ser 325 330 335
- Arg Ile Arg Pro Gly Ser Ile Ala Ala Glu Asp Val Leu His Asp Ile 340 345 350
- Gly Val Ile Ala Met Thr Ser Ser Asp Ser Gln Ala Met Gly Arg Ala 355 360 365
- Gly Glu Val Ile Pro Arg Thr Trp Gln Thr Ala Asp Lys Asn Lys Lys 370 375 380
- Glu Phe Gly Lys Leu Pro Glu Asp Gly Ala Asp Asn Asp Asn Phe Arg 385 390 395 400
- Ile Lys Arg Tyr Ile Ser Lys Tyr Thr Ile Asn Pro Ala Leu Thr His
 405 410 415
- Gly Val Ser Glu Tyr Ile Gly Ser Val Glu Glu Gly Lys Ile Ala Asp 420 425 430
- Leu Val Val Trp Asn Pro Ala Phe Phe Gly Val Lys Pro Lys Ile Val 435 440 445
- Ile Lys Gly Gly Met Val Val Phe Ser Glu Met Gly Asp Ser Asn Ala 450 455 460
- Ser Val Pro Thr Pro Gln Pro Val Tyr Tyr Arg Glu Met Phe Gly His 465 470 475 480
- His Gly Lys Ala Lys Phe Asp Thr Ser Ile Thr Phe Val Ser Lys Val
 485 490 495
- Ala Tyr Glu Asn Gly Val Lys Glu Lys Leu Gly Leu Glu Arg Lys Val
 500 505 510
- Leu Pro Val Lys Asn Cys Arg Asn Ile Thr Lys Lys Asp Phe Lys Phe 515 520 525
- Asn Asp Lys Thr Ala Lys Ile Thr Val Asp Pro Lys Thr Phe Glu Val 530 535 540
- Phe Val Asp Gly Lys Leu Cys Thr Ser Lys Pro Thr Ser Glu Val Pro





SEOIL

Leu Ala Gln Arg Tyr Thr Phe Phe 565

Printed:10-07-2001

	•	
·		
·		
	£ .	
	I .	
	·	
		•